

Summatory Multiplicative Arithmetic Functions: Scaling and Renormalization

Leonid G. Fel

Department of Civil Engineering, Technion, Haifa 32000, Israel

Abstract

We consider a wide class of summatory functions $F\{f; N, p^m\} = \sum_{k \leq N} f(p^m k)$, $m \in \mathbb{Z}_+ \cup \{0\}$, associated with the multiplicative arithmetic functions f of a scaled variable $k \in \mathbb{Z}_+$, where p is a prime number. Assuming an asymptotic behavior of summatory function, $F\{f; N, 1\} \stackrel{N \rightarrow \infty}{\sim} G_1(N) [1 + \mathcal{O}(G_2(N))]$, where $G_1(N) = N^{a_1} (\log N)^{b_1}$, $G_2(N) = N^{-a_2} (\log N)^{-b_2}$ and $a_1, a_2 \geq 0$, $-\infty < b_1, b_2 < \infty$, we calculate a renormalization function defined as a ratio, $R(f; N, p^m) = F\{f; N, p^m\} / F\{f; N, 1\}$, and find its asymptotics $R_\infty(f; p^m)$ when $N \rightarrow \infty$. We prove that the renormalization function is multiplicative, i.e., $R_\infty(f; \prod_{i=1}^n p_i^{m_i}) = \prod_{i=1}^n R_\infty(f; p_i^{m_i})$ with n distinct primes p_i . We extend these results on the others summatory functions $\sum_{k \leq N} f(p^m k^l)$, $m, l, k \in \mathbb{Z}_+$ and $\sum_{k \leq N} \prod_{i=1}^n f_i(k p^{m_i})$, $f_i \neq f_j$, $m_i \neq m_j$. We apply the derived formulas to a large number of basic summatory functions including the Euler $\phi(k)$ and Dedekind $\psi(k)$ totient functions, divisor $\sigma_n(k)$ and prime divisor $\beta(k)$ functions, the Ramanujan sum $C_q(n)$ and Ramanujan τ Dirichlet series, and others.

Keywords: Multiplicative number theory, Summatory functions, Asymptotic analysis.

2000 Mathematics Subject Classification: 11N37, 11N56

CONTENTS

1	Summatory Multiplicative Functions with Scaled Variable	2
1.1	Asymptotic Growth of Summatory Functions	4
1.2	Scaling Equation for Summatory Functions	8
2	Renormalization Function with Simple Scaling	10
2.1	Convergence of $\mathcal{R}_1(f; N, p^m)$	11
2.2	Convergence of $\mathcal{R}_2(f; N, p^m)$	14

2.3	Rational Representation of Renormalization Function	17
3	Multiplicativity of Renormalization Function with Complex Scaling	19
3.1	Convergence of $\mathcal{R}_1(f; N, \prod_{i=1}^n p_i^{m_i})$	20
3.2	Convergence of $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$	22
3.3	Asymptotics of Summatory Functions $\sum_{k_1, k_2 \leq N} f(k_1 k_2)$	25
4	Renormalization of Dirichlet Series and Others Summatory Functions	26
4.1	Renormalization of Summatory Function $\sum_{k \leq N} \prod_{i=1}^n f_i(k p_i^{m_i})$	26
4.2	Renormalization of the Dirichlet Series $\sum_{k=1}^{\infty} f(k p^m) k^{-s}$	29
4.3	Renormalization of Summatory Function $\sum_{k \leq N} f(k^n p^m)$	30
5	Renormalization of the Basic Summatory Functions	32
5.1	Renormalization of Summatory Totient Functions	32
5.1.1	Euler $\phi(k)$ function	33
5.1.2	Jordan $J_v(k)$ and Dedekind $\psi(k)$ functions	33
5.2	Renormalization of Summatory Non-Totient Functions	34
5.2.1	Divisor function $\sigma_a(k)$ and prime divisor function $\beta(k)$	34
5.2.2	Piltz function $d_n(k)$ and the sum of two squares function $r_2(k)$	36
5.2.3	Ramanujan sum $C_q(n)$ and Abelian group enumeration function $\alpha(k)$	37
5.3	Renormalization of Summatories Associated with Ramanujan's τ Function	38
5.3.1	Renormalization of the Ramanujan τ Dirichlet series	40
5.3.2	Renormalization functions $R_{\infty}(\tau^2, p^m)$ and $R_{\infty}(\tau^2 \cdot k^{-25/2}, p^m)$	41
6	Numerical Verification	42

1 SUMMATORY MULTIPLICATIVE FUNCTIONS WITH SCALED VARIABLE

Among summatory arithmetic functions $\sum_{k \leq N} f(k)$, $k \in \mathbb{Z}_+$, of various $f(k)$ the most utilized are the basic multiplicative functions $f(k)$ and their algebraic combinations. Study of different summatory functions and their asymptotics has a long history [2], [9] and [21]. Their list includes

totient functions: the Euler $\phi(k)$, Dedekind $\psi(k)$ and Jordan $J_n(k)$, and non totient functions: the Möbius $\mu(k)$ and the n -order Möbius $\mu_n(k)$, Liouville $\lambda(k)$, Piltz $d_n(k)$, divisor $\sigma_n(k)$, prime divisor $\beta(k)$, non isomorphic Abelian group enumeration function $\alpha(k)$, exponentiation of additive functions $\omega(k)$ and $\Omega(k)$ which give the numbers of distinct prime dividing k and total prime factors of k counted with multiplicities, respectively. The whole family of multiplicative arithmetic functions is much wider, e.g., the number $q_n(k)$ of representations of k by sum of two integral n th powers [29], the number $r_n(k)$ of representations of k by sum of n integer squares [21], the Legendre and Zsigmondy totient functions [9] and the Nagell totient function [23], the non isomorphic solvable [18] and nilpotent [32] finite group enumeration functions, the Gauss [21], Ramanujan [21] and Kloosterman [17] sums, the Ramanujan $\tau(k)$ function [31] and others.

In this article we study a family of summatory multiplicative arithmetic functions with scaled summation variable, $F\{f; N, p^m\} = \sum_{k \leq N} f(p^m k)$, $k, m \in \mathbb{Z}_+$, where p is a prime number. For this reason we use hereafter the notation $F\{f; N, 1\}$ for unscaled summatory function. A description of asymptotics of $F\{f; N, 1\}$, $N \rightarrow \infty$, assumes that we know two characteristics, its leading and error terms, $G_1(N)$ and $G_2(N)$, i.e., $F\{f; N, 1\} \stackrel{N \rightarrow \infty}{\sim} G_1(N) [1 + \mathcal{O}(G_2(N))]$.

In section 1.1 we introduce universality classes $\mathbb{B}\{G_1(N); G_2(N)\}$ of arithmetic functions $f(k)$ such that different functions possess the same $G_1(N)$ and $G_2(N)$. By inspection of a vast number of multiplicative functions $f(k)$ we focus on their wide class, $G_1(N) = N^{a_1} (\log N)^{b_1}$, $G_2(N) = N^{-a_2} (\log N)^{-b_2}$, where $a_1, a_2 \geq 0$ and $-\infty < b_1, b_2 < \infty$. In section 1.2 we derive a functional equation defined at different scales,

$$F\{f; N, p^m\} = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r(f; p^m) F\left\{f; \left\lfloor \frac{N}{p^r} \right\rfloor, 1\right\}, \quad (1.1)$$

where the characteristic functions $L_r(f; p^m)$ are satisfied the recursive equations,

$$L_r(f; p^m) = f(p^{m+r}) - \sum_{j=0}^{r-1} L_j(f; p^m) f(p^{r-j}), \quad L_0(f; p^m) = f(p^m), \quad f(1) = 1, \quad (1.2)$$

and $\lfloor u \rfloor$ denotes the largest integer not exceeding u . The functions $L_r(f; p^m)$ are calculated in (1.15) and their behavior in r is crucial for convergence of numerical series. This is a subject of special discussion in the next section.

In section 2 we define the renormalization function $R(f; N, p^m)$ and its asymptotics,

$$R(f; N, p^m) = \frac{F\{f; N, p^m\}}{F\{f; N, 1\}}, \quad R_\infty(f, p^m) = \lim_{N \rightarrow \infty} R(f; N, p^m). \quad (1.3)$$

The aim of this paper is to study the asymptotic renormalization function $R_\infty(f, p^m)$ in various aspects: (a) its existence as the convergent numerical series, (b) its multiplicativity property

without specifying the function $f(k)$, (c) formulas for $R_\infty(f(k) \cdot k^{-s}, p^m)$ for corresponding Dirichlet series, (d) formulas for $R_\infty(f(k^n), p^m)$ for different arithmetic functions f , (e) formulas $R_\infty(f, p^m)$ for basic arithmetic function f . For short, we'll often skip the word 'asymptotic' and refer to $R_\infty(f, p^m)$ as renormalization function if this would not mislead the readers.

Imposing the constraints on $L_r(f; p^m)$, in section 2.1 we prove two Lemmas on convergence of numerical series and calculate asymptotics of renormalization functions. In section 2.2 we show that by these constraints the error term $G_2(N)$ does not contribute to $R_\infty(f, p^m)$. In section 2.3 we give a rational representation for $R_\infty(f, p^m)$ which is much easier to implement in analytic calculations.

In section 3 we prove that the renormalization function has multiplicative property in the following sense, $R_\infty(f; \prod_{i=1}^n p_i^{m_i}) = \prod_{i=1}^n R_\infty(f; p_i^{m_i})$ with n distinct primes $p_i \geq 2$. Making use of renormalization approach we also calculate the summatory functions $\sum_{k_1, k_2 \leq N} f(k_1 k_2)$.

In section 4 we extend the renormalization approach on summatory $\sum_{k \leq N} \prod_{i=1}^n f_i(k p^{m_i})$, where $f_i \neq f_j$, $m_i \neq m_j$, and the corresponding Dirichlet series $\sum_{k=1}^\infty f(k p^m) k^{-s}$, $k, n \in \mathbb{Z}_+$. We also study the renormalization of summatory function $\sum_{k \leq N} f(k^n p^m)$.

In section 5 we apply formulas, derived in section 2, to calculate $R_\infty\{f; p^m\}$ for basic multiplicative arithmetic functions f and their combinations. Almost all summatory functions are treated by Theorem 2 based on a simple calculation of $f(p^r)$ and avoiding a cumbersome calculation of characteristic functions $L_r(f; p^m)$. The renormalization functions are given by algebraic and non-algebraic expressions as well, e.g., see $R_\infty(\sigma_0^n; p^m)$ in (5.11) and $R_\infty(\sigma_1/\sigma_0; p)$ in (5.15), respectively. In this conjunction, the Ramanujan τ function is of particular interest: in contrast to many other functions $f(k)$ its value at $k = p^r$ is given by a heavy formula (5.31), while the characteristic functions $L_r(\tau; p^m)$ and $L_r(\tau^2; p^m)$ have been calculated in a simple form suitable for explicit calculation of $R_\infty(\tau \cdot k^{-s}, p^m)$ and $R_\infty(\tau^2, p^m)$. We have found a new identity (5.37) for the Ramanujan τ function.

In section 6 we give a numerical verification to renormalization approach developed in this article by numerical calculations and show its validity with high precision.

1.1 ASYMPTOTIC GROWTH OF SUMMATORY FUNCTIONS

Consider the summatory multiplicative arithmetic function $F\{f; N, 1\}$ and represent its asymptotics in N by using one constant $\mathcal{F}$ and two positive definite functions $G_1(N)$ and $G_2(N)$,

$$F\{f; N, 1\} \stackrel{N \rightarrow \infty}{\sim} \mathcal{F} G_1(N) [1 + \mathcal{O}(G_2(N))] , \quad \lim_{N \rightarrow \infty} G_2(N) = 0 , \quad G_1(N), G_2(N) > 0 , \quad (1.4)$$

where "O" stands for the "big-O" Landau symbol.

Asymptotic growth of $F\{f; N, 1\}$ is determined by its leading term and is given by nondecreasing function $G_1(N)$, i.e., either increasing or unity, while the decreasing function $G_2(N)$ stands for the error term. The constant $\mathcal{F}$ is introduced to distinguish summatories with similar functions $G_1(N)$ and $G_2(N)$, e.g., see $F\{1/\phi; N, 1\}$, $F\{1/\psi; N, 1\}$ and $F\{1/\sigma_1; N, 1\}$ in Table 1.

Different arithmetic functions $f(k)$ may possess the same $G_1(N)$ and $G_2(N)$, therefore the whole set of $f(k)$ can be decomposed into different universality classes $\mathbb{B}\{G_1(N); G_2(N)\}$ as follows,

$$f(k) \in \mathbb{B}\{G_1(N); G_2(N)\}, \quad \text{where} \quad (1.5)$$

$$\mathbb{B}\{G_1(N); G_2(N)\} = \left\{ f(k) \left| \frac{F\{f; N, 1\}}{\mathcal{F} G_1(N)} \stackrel{N \rightarrow \infty}{\rightleftharpoons} 1, \left| \frac{F\{f; N, 1\}}{\mathcal{F} G_1(N)} - 1 \right| \stackrel{N \rightarrow \infty}{<} \mathcal{C} G_2(N); 0 < \mathcal{C}, \mathcal{F} < \infty \right\}$$

Below we give examples of various multiplicative functions $f(k)$ which belong to the different universality classes,

$$\begin{aligned} \frac{1}{\phi(k)}, \frac{1}{\psi(k)}, \frac{\phi(k)}{k^2}, \frac{1}{\sigma_1(k)} &\in \mathbb{B} \left\{ \log N; \frac{1}{\log N} \right\}, \quad \frac{\phi(k)}{k}, \frac{\sigma_1(k)}{k}, \sigma_{-1}(k) \in \mathbb{B} \left\{ N; \frac{\log N}{N} \right\}, \\ \sigma_0(k), 2^{\omega(k)}, \mu^2(k) 2^{\omega(k)} &\in \mathbb{B} \left\{ N \log N; \frac{1}{\log N} \right\}, \quad \frac{\mu^2(k)}{k}, \alpha(k), \beta(k) \in \mathbb{B} \left\{ N; \frac{1}{\sqrt{N}} \right\}. \end{aligned}$$

By inspection of a vast number of multiplicative functions $f(k)$ with known asymptotics $G_1(N)$ and $G_2(N)$ (see Tables 1 and 2) in this paper we focus on their most wide class,

$$\begin{aligned} G_1(N) &= N^{a_1} (\log N)^{b_1}, & \begin{cases} a_1 > 0, & -\infty < b_1 < \infty, \\ a_1 = 0, & 0 \leq b_1 < \infty, \end{cases} \\ G_2(N) &= \frac{1}{N^{a_2} (\log N)^{b_2}}, & \begin{cases} a_2 > 0, & -\infty < b_2 < \infty, \\ a_2 = 0, & 0 < b_2 < \infty. \end{cases} \end{aligned} \quad (1.6)$$

Remark 1 Tables 1 and 2 do not present any example of multiplicative arithmetic functions $f(k)$ of the special universality classes $\mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$ such that

$$0 < a_1 < a_2, -\infty < b_1, b_2 < \infty \quad \text{and} \quad a_1 = 0 < a_2, 0 \leq b_1 < b_2 < \infty. \quad (1.7)$$

Despite an extensive search in the available literature we have not found such functions there.

In Tables 1 and 2 we present a long list of multiplicative functions which belong to one of the universality classes $\mathbb{B}\{G_1(N); G_2(N)\}$ satisfying (1.6). We use the standard notations for the functions mentioned in section 1. Here $\zeta(s)$ stands for the Riemann zeta function, the explicit expressions for A_n , B_s , D_n , E_n , K_n and I_n and values for C_i are given in the corresponding references. The values of C_2 , C_8 and C_{10} were calculated by author and marked by ($\star$). The error terms for summatories of $|\tau(k)|$, $\tau^2(k)$, $\tau^4(k)$ and $\tau^2(k)/\sqrt{k^{25}}$ are unknown to date.

Table 1. Summatory multiplicative arithmetic functions and their asymptotics.

$f(k)$	$\mathcal{F}$	$G_1(N)$	$G_2(N)$	Ref
$\mu^2(k)$	$\zeta^{-1}(2)$	N	$N^{-1/2}$	[5]
$\mu^2(k)/\phi(k)$	1	$\log N$	$\log^{-1} N$	[47]
$\mu_n(k)$	$A_n, n \geq 2$	N	$N^{\frac{1}{n}-1} \log N$	[37], p.193
$1/\phi(k)$	$\zeta(2)\zeta(3)/\zeta(6)$	$\log N$	$\log^{-1} N$	[30]
$k/\phi(k)$	$\zeta(2)\zeta(3)/\zeta(6)$	N	$N^{-1} \log N$	[41]
$\psi(k)$	$1/2 \zeta(2)/\zeta(4)$	N^2	$N^{-1} \log N$	[2], p.72
$1/\psi(k)$	$C_1 \simeq 0.37396$	$\log N$	$\log^{-1} N$	[40]
$\sigma_{-b}(k)$	$\zeta(b+1)$	$N, b > 0, b \neq 1$	$N^{-\min\{1,b\}}$	[2], p.61
$\sigma_{-1}(k)$	$\zeta(2)$	N	$N^{-1} \log N$	[2], p.61
$\sigma_a(k)$	$\zeta(a+1)/(a+1)$	$N^{a+1}, a > 0, a \neq 1$	$N^{-\min\{1,a\}}$	[2], p.60
$\sigma_1(k)$	$\zeta(2)/2$	N^2	$N^{-1} \log^{2/3} N$	[46]
$\sigma_a^2(k)$	$\frac{\zeta^2(1+ a)\zeta(1+2 a)}{\zeta(2+2 a)}$	$N^{1+a+ a }, 0 < a < 1$	$N^{- a } \log N$	[4]
$\sigma_1^2(k)$	$5/6 \zeta(3)$	N^3	$N^{-1} \log^{5/3} N$	[42]
$\sigma_0^2(k)$	$D_n, D_1=1, D_2=\pi^{-2}$	$N \cdot \log^{2^n-1} N$	$\log^{-1} N$	[34], [48]
$1/\sigma_0(k)$	$D_{-1} \simeq 0.5469$	$N \cdot \log^{-1/2} N$	$\log^{-1} N$	[34], [48]
$\sigma_1(k)/\phi(k)$	$C_2 \simeq 3.6174 \quad \star$	N	$N^{-1} \log^2 N$	[4]
$\sigma_1(k)/\sigma_0(k)$	$C_3 \simeq 0.3569$	$N^2 \log^{-1/2} N$	$\log^{-1} N$	[3]
$1/\sigma_1(k)$	$C_4 \simeq 0.6728$	$\log N$	$\log^{-1} N$	[40]
$d_n(k)$	$1/\Gamma(n)$	$N \cdot \log^{n-1} N$	$\log^{-1} N$	[20], [39]
$d_n^2(k)$	$E_n, E_2 = D_2$	$N \cdot \log^{n^2-1} N$	$\log^{-1} N$	[25]
$1/d_n(k)$	$K_n, K_2 = D_{-1}$	$N \cdot \log^{1/n-1} N$	$\log^{-1} N$	[24]
$\beta(k)$	$\zeta(2)\zeta(3)/\zeta(6)$	N	$N^{-1/2}$	[26]
$\alpha(k)$	$\prod_{l=2}^{\infty} \zeta(l) \simeq 2.29486$	N	$N^{-1/2}$	[13]
$1/\alpha(k)$	$C_5 \simeq 0.75204$	N	$N^{-1/2} \log^{-1/2} N$	[28], p.16
$2^{\omega(k)}$	$\zeta^{-1}(2)$	$N \cdot \log N$	$\log^{-1} N$	[39]
$3^{\omega(k)}$	$C_6 \simeq 0.14338$	$N \cdot \log^2 N$	$\log^{-1} N$	[45], p.53
$2^{\Omega(k)}$	$C_7 \simeq 0.27317$	$N \cdot \log^2 N$	$\log^{-1} N$	[19]
$q_n(k)$	$2\Gamma^2(n^{-1})/(n\Gamma(2n^{-1}))$	$N^{2/n}, n \geq 3$	$N^{-1/n(1+1/n)}$	[29]. p.143
$r_2(k)=q_2(k)$	π	N	$N^{-1/2}$	[21]
$ \tau(k) $	$C_8 \simeq 0.0996 \quad \star$	$N^{13/2} \log^{-1+8/(3\pi)} N$	?	[11], [35]
$\tau^2(k)$	$C_9 \simeq 0.032007$	N^{12}	?	[31]
$\tau^4(k)$	$C_{10} \simeq 0.0026 \quad \star$	$N^{23} \log N$	?	[35]

Table 2. Other summatory functions and the Dirichlet series of multiplicative functions.

$f(k)$	$\mathcal{F}$	$G_1(N)$	$G_2(N)$	Ref
$\mu(k)/k^s$	$\zeta^{-1}(s), s > 1$	N^0	$N^{-(s-1)}$	[2], p.231
$\mu^2(k)/k^s$	$\zeta(s)/\zeta(2s), s > 1$	N^0	$N^{-(s-1)}$	[2], p.241
$\mu^2(k)/k$	$\zeta^{-1}(2)$	$\log N$	$\log^{-1} N$	[38], p.195
$\lambda(k)/k^s$	$\zeta(2s)/\zeta(s), s > 1$	N^0	$N^{-(s-1)}$	[2], p.231
$\phi(k)/k^s$	$\zeta^{-1}(2)/(2-s), 0 < s \leq 1$	N^{2-s}	$N^{-1} \log N$	[2], p.71
$\phi(k)/k^s$	$\zeta^{-1}(2)/(2-s), 1 < s < 2$	N^{2-s}	$N^{-(2-s)}$	[2], p.71
$\phi(k)/k^2$	$\zeta^{-1}(2)$	$\log N$	$\log^{-1} N$	[2], p.71
$\phi(k)/k^s$	$\zeta(s-1)/\zeta(s), s > 2$	N^0	$N^{-(s-2)}$	[2], p.231
$(\phi(k)/k)^s$	$B_s, B_1 = \zeta^{-1}(2)$	$N, s > 0$	$N^{-1} \log^s N$	[6]
$\sigma_0(k)/k^s$	$(1-s)^{-1}, 0 < s < 1$	$N^{1-s} \log N$	$\log^{-1} N$	[2], p.70
$\sigma_0(k)/k$	$1/2$	$\log^2 N$	$\log^{-1} N$	[2], p.70
$\sigma_0(k)/k^s$	$\zeta^2(s), s > 1$	N^0	$N^{-(s-1)} \log N$	[2], p.231
$(\sigma_1(k)/k)^s$	$I_s, I_1 = \zeta(2), I_2 = \frac{5}{2}\zeta(3)$	$N, s > 0$	$N^{-1} \log^s N$	[42], [4]
$r_2(k)/k$	π	$\log N$	$\log^{-1} N$	[21]
$\tau^2(k)/k^{25/2}$	$C_{11} \simeq 1.58824$	N^0	?	[10]

A relationship between the multiplicative properties of arithmetic functions and asymptotics of their summatory functions is not straightforward. In other words, a correspondence:

$$f(k) \text{ is a multiplicative function} \longleftrightarrow f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\} \quad (1.8)$$

is neither bijective nor injective. Indeed, the direction ‘ $\longleftarrow$ ’ is not holding since there exists a non multiplicative function $f(k) = \log \phi(k) / \log \sigma_1(k)$ which has the summatory function $F\{f; N, 1\} \stackrel{N \rightarrow \infty}{\sim} N + \mathcal{O}(N \log^{-1} N)$ [28].

Regarding another direction ‘ $\longrightarrow$ ’ there exist multiplicative arithmetic functions with summatory growth that differs from $N^{a_1} (\log N)^{b_1}$ and come by enumeration of finite groups. Let $\chi(k)$ be a number of nilpotent groups of order k which is multiplicative, because each finite nilpotent group is a direct product of its Sylow subgroups [32]. When $k = p^r$, for a prime p , it is known [22] that $\chi(k) \simeq p^{(2/27+o(1))r^3}$, i.e., of the order $k^{(\log_p k)^2}$. Consider its summatory $F[\chi; N, 1]$ which is the number of nilpotent groups of order at most N . For some r we have $2^r \leq N < 2^{r+1}$.

Then for some $0 < s < 1$ we have,

$$F[\chi; N, 1] \geq \chi(2^r) = 2^{(2/27+o(1))r^3} \geq (2^{r+1})^{2s/27(r+1)^2} \rightarrow F[\chi; N, 1] \geq N^{2s/27(\log N)^2}, \quad (1.9)$$

and therefore, $\chi(k) \notin \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$. Note that the number $\chi(k, c, d)$ of nilpotent groups of order k , of nilpotency class at most n , generated by at most m elements does belong to the universality class defined in (1.6) where a_i and b_i depend on n and m [36].

Other examples of multiplicative function with summatory growth $N^{a_1} (\log N)^{b_1}$ and the error term, which is different than $N^{-a_2} (\log N)^{-b_2}$, were given in [46],

$$\phi(k) \in \mathbb{B} \left(N^2; \frac{1}{N} (\log N)^{2/3} (\log \log N)^{4/3} \right), \quad \frac{\phi(k)}{k} \in \mathbb{B} \left(N; \frac{1}{N} (\log N)^{2/3} (\log \log N)^{4/3} \right).$$

They can also be encompassed within the universality classes by extending the latter on much wider family of asymptotics, e.g., $\mathbb{B} \left\{ N^{a_1} (\log N)^{b_1} (\log \log N)^{c_1}; N^{-a_2} (\log N)^{-b_2} (\log \log N)^{-c_2} \right\}$. Keeping in mind such option, we continue to study the summatory multiplicative functions with universality classes of asymptotics given in (1.6).

1.2 SCALING EQUATION FOR SUMMATORY FUNCTIONS

Represent $F\{f; N, p^m\}$ as a sum of two summatory functions

$$F\{f; N, p^m\} = \sum_{\substack{k=1 \\ p \nmid k}}^N f(p^m k) + \sum_{\substack{k=p \\ p \mid k}}^N f(p^m k). \quad (1.10)$$

Making use of multiplicativity, $f(k_1 k_2) = f(k_1) f(k_2)$ if $\gcd(k_1, k_2) = 1$, $f(1) = 1$, we get

$$\begin{aligned} F\{f; N, p^m\} &= f(p^m) \sum_{\substack{k=1 \\ p \nmid k}}^N f(k) + \sum_{\substack{k=p \\ p \mid k}}^N f(p^m k) = f(p^m) \left(\sum_{k=1}^N f(k) - \sum_{\substack{k=p \\ p \mid k}}^N f(k) \right) + \sum_{\substack{k=p \\ p \mid k}}^N f(p^m k) \\ &= f(p^m) F\{f; N, 1\} - f(p^m) \sum_{l=1}^{N_1} f(pl) + \sum_{l=1}^{N_1} f(p^{m+1} l) \\ &= f(p^m) F\{f; N, 1\} - f(p^m) F\{f; N_1, p\} + F\{f; N_1, p^{m+1}\}. \end{aligned} \quad (1.11)$$

where $N_r = \lfloor N/p^r \rfloor$. The recursion (1.11) holds for any N_r , i.e.,

$$F\{f; N_r, p^m\} = f(p^m) F\{f; N_r, 1\} - f(p^m) F\{f; N_{r+1}, p\} + F\{f; N_{r+1}, p^{m+1}\}. \quad (1.12)$$

Substituting (1.12) into (1.11) we obtain

$$\begin{aligned} F\{f; N, p^m\} &= f(p^m) F\{f; N, 1\} - f(p^m) [f(p) F\{f; N_1, 1\} - f(p) F\{f; N_2, p\} + F\{f; N_2, p^2\}] \\ &\quad + f(p^{m+1}) F\{f; N_1, 1\} - f(p^{m+1}) F\{f; N_2, p\} + F\{f; N_2, p^{m+2}\} \\ &= f(p^m) F\{f; N, 1\} + [f(p^{m+1}) - f(p^m) f(p)] F\{f; N_1, 1\} \\ &\quad + [f(p^m) f(p) - f(p^{m+1})] F\{f; N_2, p\} - f(p^m) F\{f; N_2, p^2\} + F\{f; N_2, p^{m+2}\}. \end{aligned}$$

Continuing this procedure recursively we get finally,

$$F\{f; N, p^m\} = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r(f; p^m) F\{f; N_r, 1\}, \quad \text{where} \quad (1.13)$$

$$L_r(f; p^m) = f(p^{m+r}) - \sum_{j=0}^{r-1} L_j(f; p^m) f(p^{r-j}), \quad L_0(f; p^m) = f(p^m). \quad (1.14)$$

The straightforward calculations of $L_r(f; p^m)$ give

$$L_1(f; p^m) = f(p^{m+1}) - f(p^m) f(p), \quad (1.15)$$

$$L_2(f; p^m) = f(p^{m+2}) - f(p^m) f(p^2) - f(p^{m+1}) f(p) + f(p^m) f^2(p),$$

$$L_3(f; p^m) = f(p^{m+3}) - f(p^m) f(p^3) - f(p^{m+1}) f(p^2) + 2f(p^m) f(p^2) f(p) - f(p^{m+2}) f(p) + f(p^{m+1}) f^2(p) - f(p^m) f^3(p), \quad \text{etc},$$

such that for $p = 1$ or $m = 0$ we have, $L_0(f; 1) = 1$ and $L_r(f; 1) = 0$, $r \geq 1$.

By (1.14) or (1.15) the general formulas for $L_r(f; p^m)$ can be calculated by induction for simple arithmetic functions $f(k)$ and $1/f(k)$ such that $f(p^m) = A_f p^{m-1}$, $m \geq 1$,

$$L_r(f; p^m) = A_f(p - A_f)^r p^{m-1}, \quad L_r\left(\frac{1}{f}; p^m\right) = \frac{1}{A_f} \left(\frac{1}{p} - \frac{1}{A_f}\right)^r \frac{1}{p^{m-1}}, \quad r \geq 1, \quad (1.16)$$

and A_f denotes a real constant, e.g., $A_\phi = p - 1$, $A_\psi = p + 1$ that gives

$$\begin{cases} L_r(\phi; p^m) = p^{m-1}(p - 1), \\ L_r(\psi; p^m) = (-1)^r p^{m-1}(p + 1), \end{cases} \quad L_r\left(\frac{1}{\phi}; p^m\right) = \frac{(-1)^r p^{1-r-m}}{(p-1)^{r+1}}, \quad L_r\left(\frac{1}{\psi}; p^m\right) = \frac{p^{1-r-m}}{(p+1)^{r+1}}.$$

Another example of arithmetic functions leading to $L_r(f; p^m) = 0$, $r \geq 1$, is that when $A_f = p$ in (1.16) or $f(p^m) = c^m$, e.g., $2^{\Omega(p^m)} = 2^m$ and $\lambda(p^m) = (-1)^m$.

In general case of $f(k)$ the formula of $L_r(f; p^m)$ with arbitrary $r \geq 0$ can be hardly recognized by its partial expressions, e.g., for $f(k) = 1/\sigma_0(k)$,

$$L_0 = \frac{1}{m+1}, \quad L_1 = \frac{m m!}{2(m+2)!}, \quad L_2 = \frac{(5m+7) m m!}{12(m+3)!}, \quad L_3 = \frac{(9m^2 + 35m + 32) m m!}{24(m+4)!}. \quad (1.17)$$

In section 5.2.1, formula (5.13), we show that L_r in (1.17) come as coefficients in the series expansion of the function involving logarithmic and hypergeometric functions.

Remark 2 Consider an integer N in the range $p^{\bar{r}} \leq N < p^{\bar{r}+1}$ where $\bar{r} = \lfloor \log_p N \rfloor$ and write (1.13) as follows,

$$F\{f; N, p^m\} = \sum_{r=0}^{\bar{r}-1} L_r(f; p^m) F\{f; N_r, 1\} + L_{\bar{r}}(f; p^m) F\{f; N_{\bar{r}}, 1\}. \quad (1.18)$$

where $1 \leq N_{\bar{r}} = \lfloor N/p^{\bar{r}} \rfloor < p$ and $F[f; N_{\bar{r}}, 1] = \sum_{k=1}^{N_{\bar{r}}} f(k) < \sum_{k=1}^p f(k)$ is a finite number.

We make use of representation (1.18) in section 2.1 when studying the asymptotics of renormalization functions for universality classes with $b_1 < 0$ (Lemma 2) and $b_1 < b_2$ (Lemma 4).

2 RENORMALIZATION FUNCTION WITH SIMPLE SCALING

Define the renormalization function as a ratio of two summatory functions

$$R(f; N, p^m) = \frac{F\{f; N, p^m\}}{F\{f; N, 1\}}, \quad R(f; N, 1) = 1. \quad (2.1)$$

Substituting (1.4) into (2.1) we get its asymptotic behavior

$$R(f; N, p^m) \stackrel{N \rightarrow \infty}{\equiv} \frac{\mathcal{R}_1(f; N, p^m)}{1 + \mathcal{O}(G_2(N))} + \frac{\mathcal{R}_2(f; N, p^m)}{1 + \mathcal{O}(G_2(N))},$$

where $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ are defined according to (1.13) as follows

$$\mathcal{R}_1(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r(f; p^m) \frac{G_1(N_r)}{G_1(N)}, \quad (2.2)$$

$$\mathcal{R}_2(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r(f; p^m) \frac{G_1(N_r)}{G_1(N)} \mathcal{O}(G_2(N_r)). \quad (2.3)$$

If both numerical series $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ converge when $N \rightarrow \infty$, then

$$\lim_{N \rightarrow \infty} R(f; N, p^m) := R_\infty(f; p^m) = \lim_{N \rightarrow \infty} \mathcal{R}_1(f; N, p^m) + \lim_{N \rightarrow \infty} \mathcal{R}_2(f; N, p^m). \quad (2.4)$$

What can be said about convergence of $R(f; N, p^m)$ without knowing exactly the multiplicative function $f(k)$ itself? Formulas (2.2) and (2.3) for $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ indicate that a large portion of information is hidden in the asymptotics $G_1(N)$ and $G_2(N)$.

Substitute $G_1(N) = N^{a_1} (\log N)^{b_1}$ into (2.2) and get

$$\mathcal{R}_1(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{L_r(f; p^m)}{p^{a_1 r}} \left(1 - \frac{r}{\log_p N}\right)^{b_1}. \quad (2.5)$$

Regarding $\mathcal{R}_2(f; N, p^m)$, which is responsible for contribution of the error term into $R(f; N, p^m)$, note that according to the definition (1.5) we get $\mathcal{O}(G_2(N)) \leq \mathcal{C} G_2(N)$. Applying this inequality to formula (2.3),

$$|\mathcal{R}_2(f; N, p^m)| \leq \mathcal{C} \sum_{r=0}^{\lfloor \log_p N \rfloor} |L_r(f; p^m)| \frac{G_1(N_r)}{G_1(N)} G_2(N_r),$$

and substituting $G_2(N) = N^{-a_2} (\log N)^{-b_2}$ into the last expression we get an estimate,

$$|\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{C}}{N^{a_2} (\log N)^{b_2}} \mathcal{R}_3(f; N, p^m), \quad \text{where} \quad (2.6)$$

$$\mathcal{R}_3(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{|L_r(f; p^m)|}{p^{r(a_1 - a_2)}} \left(1 - \frac{r}{\log_p N}\right)^{b_1 - b_2}. \quad (2.7)$$

In sections 2.1 and 2.2 we give a detailed analysis of convergence of $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ for multiplicative functions $f(k)$ of several universality classes. Start with a specific class of $f(k)$ and, assuming only $L_r(f; p^m) \geq 0$, prove that the convergence of $\mathcal{R}_1(f; N, p^m)$ implies the convergence of $\mathcal{R}_2(f; N, p^m)$ to zero.

Proposition 1 *Let $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_1} (\log N)^{-b_2} \right\}$ be given such that $L_r(f; p^m) \geq 0$ and let $\mathcal{R}_1(f; N, p^m)$ be convergent. Then $\mathcal{R}_2(f; N, p^m) \xrightarrow{N \rightarrow \infty} 0$, in each of the cases,*

- 1) $a_1 = a_2 = 0, b_1 \geq b_2 > 0,$ 2) $0 < a_2 < a_1, 0 \leq b_2 \leq b_1$ or $0 = a_2 < a_1, 0 < b_2 \leq b_1,$
- 3) $0 < a_2 < a_1, b_1 < b_2, b_1 < 0$ or $0 = a_2 < a_1, b_1 < 0 < b_2.$

Proof Keeping in mind $L_r(f; p^m) \geq 0$ and comparing (2.5) and (2.7) we conclude that if $\mathcal{R}_1(f; N, p^m)$ is convergent when $a_1 = 0$ and $b_1 \geq 0$ then $\mathcal{R}_3(f; N, p^m)$ is also convergent when $a_1 = a_2 = 0$ and $b_1 \geq b_2$. Substituting this into (2.6) we get

$$|\mathcal{R}_2(f; N, p^m)| \leq C (\log N)^{-b_2} \mathcal{R}_3(f; N, p^m), \quad (2.8)$$

that proves Proposition if $b_2 > 0$. Indeed, if b_2 is positive and $\mathcal{R}_3(f; N, p^m)$ is convergent, then the right hand side (r.h.s.) in (2.8) is convergent to zero and so does $\mathcal{R}_2(f; N, p^m)$. Applying similar arguments in the two others cases (2) and (3) we prove Proposition completely. $\square$

Table 1 shows when the 1st item in Proposition 1 can be applied: this is the inverse Dedekind function: $a_1 = a_2 = 0, b_1 = b_2 = 1$ and by (1.16) $L_r(1/\psi; p^m) \geq 0$. But neither the Euler totient function nor its inverse can be studied by Proposition 1 which is quite weak statement.

The convergence of $\mathcal{R}_1(f; N, p^m)$ implies a zero limit of $\mathcal{R}_2(f; N, p^m)$, $N \rightarrow \infty$, in much wider range of varying degrees a_1, b_1 and a_2, b_2 . Indeed, to provide the convergence of $\mathcal{R}_2(f; N, p^m)$ to zero there is no need to require the convergence of $\mathcal{R}_3(f; N, p^m)$ in (2.6) but rather to allow a growth of $\mathcal{R}_3(f; N, p^m)$ with a rate less than $N^{a_2} (\log N)^{b_2}$. However, this would require more assumptions about $L_r(f; p^m)$. In the next sections 2.1 and 2.2 we study the convergence problem in more details and prove the main result of this section in Theorem 1.

2.1 CONVERGENCE OF $\mathcal{R}_1(f; N, p^m)$

In this section we study the convergence of $\mathcal{R}_1(f; N, p^m)$ when $N \rightarrow \infty$. The conditions imposed on $L_r(f; p^m)$ provide convergence of $\mathcal{R}_1(f; N, p^m)$. Throughout this and the next sections we repeatedly make use of the squeeze (SQ) theorem [43], which is also known as the pinching or sandwich theorems.

Lemma 1 *Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1} ; G_2(N) \right\}$, $a_1 \geq 0$, $b_1 \geq 0$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$. Then*

$$\lim_{N \rightarrow \infty} \mathcal{R}_1(f; N, p^m) = \sum_{r=0}^{\infty} \frac{L_r(f; p^m)}{p^{a_1 r}}. \quad (2.9)$$

Proof First, consider $\mathcal{R}_1(f; N, p^m)$ given in (2.5) when $a_1 \geq 0$, $b_1 \in \mathbb{Z}_+ \cup \{0\}$, i.e., b_1 is a nonnegative integer. After binomial expansion in (2.5) we get,

$$\mathcal{R}_1(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{L_r(f; p^m)}{p^{a_1 r}} + \sum_{k=1}^{b_1} \left(\frac{-1}{\log_p N} \right)^k \binom{b_1}{k} \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{L_r(f; p^m)}{p^{a_1 r}} r^k. \quad (2.10)$$

Focus on the inner sum

$$\mathcal{R}_4(f; N, p^m, k) = \frac{1}{(\log_p N)^k} \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{L_r(f; p^m)}{p^{a_1 r}} r^k, \quad 1 \leq k \leq b_1, \quad (2.11)$$

and prove that the sum in (2.11) is convergent absolutely. To find an estimate for $\mathcal{R}_4(f; N, p^m, k)$ we have to consider the last sum at interval $(r_*, \lfloor \log_p N \rfloor)$ where an inequality $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ holds. However, because of the prefactor $(\log_p N)^{-k}$, $k \geq 1$, a summation at interval $(0, r_* - 1)$ does not contribute to the limit when $N \rightarrow \infty$ and does not change the convergence of the entire sum (2.11). Then

$$|\mathcal{R}_4(f; N, p^m, k)| \leq \frac{1}{(\log_p N)^k} \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{|L_r(f; p^m)|}{p^{a_1 r}} r^k \leq \frac{\mathcal{K}}{(\log_p N)^k} \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{r^k}{p^{\epsilon r}}, \quad (2.12)$$

where $\epsilon = a_1 - \gamma > 0$. Denote $M = \lfloor \log_p N \rfloor$ and consider the sum in the r.h.s. of (2.12),

$$T(p, k, \epsilon, M) = \sum_{r=0}^M \frac{r^k}{p^{\epsilon r}} = \text{Li}_{-k}(p^{-\epsilon}) - p^{-\epsilon(M+1)} \Phi(p^{-\epsilon}, -k, M+1), \quad (2.13)$$

where $\text{Li}_s(z)$ and $\Phi(z, s, a)$ are the polylogarithm function and the Hurwitz-Lerch zeta function [12], respectively,

$$\text{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}, \quad \Phi(z, s, a) = \sum_{k=0}^{\infty} \frac{z^k}{(a+k)^s}. \quad (2.14)$$

Keeping in mind the asymptotics of $\Phi(z, s, a)$, $a \rightarrow \infty$, for fixed s and z (Thm.1, [15]), $\Phi(z, s, a) \simeq a^{-s}/(1-z)$, and combining it with (2.11) and (2.13), we get

$$|\mathcal{R}_4(f; N, p^m, k)| \stackrel{N \rightarrow \infty}{\leq} \mathcal{K} \left[\frac{\text{Li}_{-k}(p^{-\epsilon})}{(\log_p N)^k} - \frac{N^{-\epsilon}}{p^{\epsilon} - 1} \right], \quad 1 \leq k \leq b_1. \quad (2.15)$$

Since the polylogarithm $\text{Li}_{-k}(z)$, $0 \leq k < \infty$, is a bounded rational function in z when $0 \leq z < 1$, then for $\epsilon > 0$ the function $\text{Li}_{-k}(p^{-\epsilon})$ is also bounded. Thus, $\mathcal{R}_4(f; N, p^m, k)$ is convergent to zero and by (2.10) and (2.11) the limit (2.9) holds.

Extend this result on the entire set of the nonnegative real numbers b_1 . In order to do this, we make use of the SQ theorem and start with trivial inequalities,

$$\left(1 - \frac{r}{\log_p N}\right)^{\lfloor b_1+1 \rfloor} \leq \left(1 - \frac{r}{\log_p N}\right)^{b_1} \leq \left(1 - \frac{r}{\log_p N}\right)^{\lfloor b_1 \rfloor}, \quad 0 \leq r \leq \lfloor \log_p N \rfloor, \quad (2.16)$$

which due to (2.5) implies the following relations,

$$\mathcal{J}_1(f; p^m, \log_p N, \lfloor b_1 + 1 \rfloor) \leq \mathcal{J}_1(f; p^m, \log_p N, b_1) \leq \mathcal{J}_1(f; p^m, \log_p N, \lfloor b_1 \rfloor), \quad (2.17)$$

$$\text{where} \quad \mathcal{J}_1(f; p^m, M, b) = \sum_{r=0}^{\lfloor M \rfloor} |L_r(f; p^m)| p^{-a_1 r} \left(1 - \frac{r}{M}\right)^b, \quad b \geq 0.$$

By proof on convergence of $\mathcal{R}_1(f; N, p^m)$ with nonnegative integer degrees b_1 and by (2.17) and by the SQ theorem it follows the convergence of $\mathcal{R}_1(f; N, p^m)$ with real $b_1 \geq 0$. $\square$

Lemma 1 can be applied to the totient functions $\phi(k)$, $\psi(k)$ and their inverse $1/\phi(k)$, $1/\psi(k)$ with the functions $L_r(f; p^m)$ calculated in (1.16).

Lemma 2 *Let a function $f(k) \in \mathbb{B}\{N^{a_1}(\log N)^{b_1}; G_2(N)\}$, $a_1 > 0$, $b_1 < 0$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K}p^{\gamma r}$ for all $r \geq r_*$. Then (2.9) holds.*

Proof Consider $\mathcal{R}_1(f; N, p^m)$ given in (2.5) when $a_1 > 0$, $b_1 \in \mathbb{Z}_-$, i.e., b_1 is a negative integer. In order to avoid its divergence at $r = \log_p N$ we use the representation (1.18) in Remark 2,

$$\mathcal{R}_1(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{L_r(f; p^m)}{p^{a_1 r}} \left(1 - \frac{r}{\log_p N}\right)^{-|b_1|} + \frac{A (\log_p N)^{|b_1|}}{\mathcal{F} N^{a_1}}, \quad (2.18)$$

where $A = L_{\bar{r}}(f; p^m) F[f; N_{\bar{r}}, 1] < \infty$ and $\bar{r} = \lfloor \log_p N \rfloor$ were defined in (1.18).

Estimate $\mathcal{R}_1(f; N, p^m)$ when $|L_r(f; p^m)| < \mathcal{K}p^{\gamma r}$, $\mathcal{K} > 0$ and $\gamma < a_1$. Note that the last term in (2.18) does not contribute to the asymptotics of $\mathcal{R}_1(f; N, p^m)$ when $N \rightarrow \infty$ and therefore it can be skipped hereafter. We use an identity

$$(1-x)^{-b} = 1 + x \sum_{k=1}^b (1-x)^{-k}, \quad b \in \mathbb{Z}_+, \quad (2.19)$$

and represent (2.18) as follows,

$$\mathcal{R}_1(f; N, p^m) \simeq \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{L_r(f; p^m)}{p^{a_1 r}} + \sum_{k=1}^{|b_1|} \mathcal{R}_5(f; N, p^m, k), \quad \text{where} \quad (2.20)$$

$$\mathcal{R}_5(f; N, p^m, k) = \frac{1}{\log_p N} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{L_r(f; p^m)}{p^{a_1 r}} r \left(1 - \frac{r}{\log_p N}\right)^{-k}. \quad (2.21)$$

Note that the following inequality holds,

$$\left(1 - \frac{r}{\log_p N}\right)^{-1} \leq 1 + r \quad \text{if } 0 \leq r < \lfloor \log_p N \rfloor. \quad (2.22)$$

Substitute (2.22) into (2.21) and keep in mind that due to the prefactor $(\log_p N)^{-1}$ in the r.h.s. of Eq. (2.21) the same convergence of $\mathcal{R}_5(f; N, p^m, k)$ holds at intervals $(r_*, \lfloor \log_p N \rfloor - 1)$ and $(0, \lfloor \log_p N \rfloor - 1)$ (see discussion in proof of Lemma 1). Then we arrive at estimate,

$$|\mathcal{R}_5(f; N, p^m, k)| \leq \frac{\mathcal{K}}{\log_p N} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{r(1+r)^k}{p^{\epsilon r}} = \frac{\mathcal{K}}{\log_p N} \sum_{j=0}^k \binom{k}{j} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{r^{j+1}}{p^{\epsilon r}}.$$

The rest of the proof follows by applying the same arguments of asymptotics of the Hurwitz-Lerch zeta function, as it was done in Lemma 1,

$$|\mathcal{R}_5(f; N, p^m, k)| \leq \mathcal{K} \sum_{j=0}^k \binom{k}{j} \left[\frac{\text{Li}_{-(j+1)}(p^{-\epsilon})}{\log_p N} - \frac{1}{p^\epsilon - 1} \frac{(\log_p N)^j}{N^\epsilon} \right]. \quad (2.23)$$

By comparison the r.h.s. in (2.23) and (2.15) we conclude that $\mathcal{R}_5(f; N, p^m, k)$ is convergent to zero when $N \rightarrow \infty$. Thus, by (2.20) the limit (2.9) holds for $b_1 \in \mathbb{Z}_-$.

We extend this result by the SQ theorem on all negative real b_1 . This can be done by inequality (2.17) for another function $\mathcal{J}_2(f; p^m, M, b)$,

$$\mathcal{J}_2(f; p^m, \log_p N, \lfloor b_1 + 1 \rfloor) \leq \mathcal{J}_2(f; p^m, \log_p N, b_1) \leq \mathcal{J}_2(f; p^m, \log_p N, \lfloor b_1 \rfloor), \quad (2.24)$$

$$\text{where} \quad \mathcal{J}_2(f; p^m, M, b) = \sum_{r=0}^{\lfloor M \rfloor - 1} \frac{|L_r(f; p^m)|}{p^{a_1 r}} \left(1 - \frac{r}{M}\right)^b, \quad b < 0.$$

By proof on convergence of $\mathcal{R}_1(f; N, p^m)$ with negative integer degrees b_1 and by (2.24) and by the SQ theorem it follows the convergence of $\mathcal{R}_1(f; N, p^m)$ with real $b_1 < 0$. $\square$

2.2 CONVERGENCE OF $\mathcal{R}_2(f; N, p^m)$

In this section we consider the convergence of $\mathcal{R}_2(f; N, p^m)$, $N \rightarrow \infty$, defined in (2.2) and responsible for contribution of the error term to $R_\infty(f; p^m)$.

Lemma 3 *Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$, $b_1 \geq b_2$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$, then*

$$\lim_{N \rightarrow \infty} \mathcal{R}_2(f; N, p^m) = 0. \quad (2.25)$$

Proof Denote $b_1 - b_2 = e$ and make use of a simple inequality, $(1 - x)^e \leq 1$ when $0 \leq x \leq 1$, $e \geq 0$. Formula (2.6) can be rewritten as follows,

$$|\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{C}}{N^{a_2} (\log N)^{b_2}} \sum_{r=0}^{\lfloor \log_p N \rfloor} \frac{|L_r(f; p^m)|}{p^{(a_1 - a_2)r}} \leq \frac{\mathcal{C}\mathcal{K}}{N^{a_2} (\log N)^{b_2}} \sum_{r=0}^{\lfloor \log_p N \rfloor} p^{\nu r}, \quad (2.26)$$

where $\nu = a_2 - a_1 + \gamma$. Keep in mind that due to the prefactor $N^{-a_2} (\log N)^{-b_2}$ in (2.26), the same convergence of the r.h.s. in (2.26) holds at intervals $(r_*, \lfloor \log_p N \rfloor)$ and $(0, \lfloor \log_p N \rfloor)$ (see discussion in proof of Lemma 1).

The further calculations are dependent on the sign of ν . If $\nu < 0$ then

$$\gamma < a_1 - a_2, \quad |\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{C}\mathcal{K}}{1 - p^\nu} \frac{N^{-a_2}}{(\log N)^{b_2}}. \quad (2.27)$$

If $\nu = 0$ then

$$\gamma = a_1 - a_2, \quad |\mathcal{R}_2(f; N, p^m)| \leq \mathcal{C}\mathcal{K} \frac{N^{\gamma - a_1}}{(\log N)^{b_2 - 1}}. \quad (2.28)$$

Finally, if $\nu > 0$ then

$$\gamma > a_1 - a_2, \quad |\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{C}\mathcal{K}}{N^{a_2} (\log N)^{b_2}} \frac{p^{\nu(\lfloor \log_p N \rfloor + 1)}}{p^\nu - 1} \simeq \frac{\mathcal{C}\mathcal{K}}{1 - p^{-\nu}} \frac{N^{\gamma - a_1}}{(\log N)^{b_2}}. \quad (2.29)$$

Require now that all r.h.s. in (2.27), (2.28) and (2.29) converge to zero when $N \rightarrow \infty$. Regarding the 1st case (2.27) this always holds because by (1.6) if $a_2 = 0$ then $b_2 > 0$, and if $a_2 > 0$ then $b_2 \geq 0$, so the r.h.s. in (2.27) is decreasing function. So, it results in requirement, $\gamma < a_1 - a_2$. In two other cases we have necessary conditions,

$$a_1 - a_2 = \gamma \leq a_1 \quad \text{if} \quad b_2 > 1, \quad \text{and} \quad a_1 - a_2 = \gamma < a_1 \quad \text{if} \quad -\infty < b_2 < \infty, \quad (2.30)$$

$$a_1 - a_2 < \gamma \leq a_1 \quad \text{if} \quad b_2 > 0, \quad \text{and} \quad a_1 - a_2 < \gamma < a_1 \quad \text{if} \quad -\infty < b_2 < \infty. \quad (2.31)$$

Summarizing the necessary conditions (2.30), (2.31) and $\gamma < a_1 - a_2$, we conclude that the numerical series $\mathcal{R}_2(f; N, p^m)$ is convergent to zero absolutely when $N \rightarrow \infty$ and irrespectively to the sign of b_2 if $\gamma < a_1$. This proves formula (2.25). $\square$

Lemma 4 Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$, $b_1 < b_2$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$, then (2.25) holds.

Proof Consider $\mathcal{R}_3(f; N, p^m)$ given in (2.7) and, according to Remark 2, rewrite it as follows,

$$\mathcal{R}_3(f; N, p^m) = \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{|L_r(f; p^m)|}{p^{(a_1 - a_2)r}} \left(1 - \frac{r}{\log_p N} \right)^{-|b_2 - b_1|} + \frac{A}{\mathcal{F}} \frac{N^{-a_1}}{(\log_p N)^{b_1}}, \quad (2.32)$$

where the upper bound in the sum is taken in order to avoid its divergence and A is defined in (2.18), more details see in proof of Lemma 2, formula (2.18). The last term in (2.32) does not contribute to asymptotics of $\mathcal{R}_2(f; N, p^m)$ when $N \rightarrow \infty$ and can be skipped. Indeed, if $a_1 \geq 0$, $b_1 > 0$ it converges to zero when $N \rightarrow \infty$; in the case $a_1 = b_1 = 0$ according to (1.6) we have $a_2 > 0$ or $a_2 = 0$, $b_2 > 0$ that again makes it irrelevant due to prefactor $N^{-a_2} (\log_p N)^{-b_2}$ in formula (2.6) for $\mathcal{R}_2(f; N, p^m)$. Apply an inequality (2.22) in the range $0 \leq r \leq \lfloor \log_p N \rfloor - 1$,

$$\left(1 - \frac{r}{\log_p N}\right)^{-(b_2-b_1)} \leq (\log_p N)^{b_2-b_1},$$

and substitute it into (2.32)

$$\mathcal{R}_3(f; N, p^m) \leq (\log_p N)^{b_2-b_1} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{|L_r(f; p^m)|}{p^{r(a_1-a_2)}}. \quad (2.33)$$

If $a_2 > 0$ we apply to (2.33) the constraints on $L_r(f; p^m)$ and substitute the result into (3.6),

$$\mathcal{R}_2(f; N, p^m) \leq \frac{\mathcal{CK}}{N^{a_2} (\log_p N)^{b_1}} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} p^{\nu r}, \quad \nu = a_2 - a_1 + \gamma. \quad (2.34)$$

By comparison (2.34) with (2.26) from Lemma 3 we obtain according to (2.27), (2.28) and (2.29)

$$|\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{CK}}{p^\nu - 1} \frac{N^{\gamma-a_1}}{(\log_c N)^{b_1}}, \quad \nu > 0, \quad (2.35)$$

$$|\mathcal{R}_2(f; N, p^m)| \leq \mathcal{CK} \frac{N^{-a_2}}{(\log_c N)^{b_1-1}}, \quad \nu = 0. \quad (2.36)$$

$$|\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{CK}}{1-p^\nu} \frac{N^{-a_2}}{(\log_c N)^{b_1}}, \quad \nu < 0. \quad (2.37)$$

Thus, by (2.35), (2.36), (2.37) and inequalities $\gamma < a_1$, $a_2 > 0$ a series $\mathcal{R}_2(f; N, p^m)$ is convergent to zero when $N \rightarrow \infty$ irrespectively to the value of b_1 .

Consider another case $a_2 = 0$ and compare formulas (2.18) and (2.32) for $b_1 < 0$ and $b_1 < b_2$, respectively. A difference in degrees, $b_1 - b_2$ and b_1 , does not break the main result of Lemma 2: only the 1st leading term in (2.20) is survived when $N \rightarrow \infty$. When we apply it to (2.32) and make use of constraint on $L_r(f; p^m)$ we get,

$$\mathcal{R}_3(f; N, p^m) \stackrel{N \rightarrow \infty}{\simeq} \sum_{r=0}^{\lfloor \log_p N \rfloor - 1} \frac{L_r(f; p^m)}{p^{a_1 r}} \leq \mathcal{K} \sum_{r=0}^{\infty} p^{-\epsilon r} = \frac{\mathcal{K}}{1-p^{-\epsilon}},$$

where $\epsilon = a_1 - \gamma > 0$. Substituting the last estimate into (3.6) we obtain,

$$|\mathcal{R}_2(f; N, p^m)| \leq \frac{\mathcal{CK}}{1-p^{-\epsilon}} \cdot (\log_p N)^{-b_2}. \quad (2.38)$$

Recall that by (1.6) the degrees of the error term satisfy: if $a_2 = 0$ than $b_2 > 0$. Thus, by (2.38) the series $\mathcal{R}_2(f; N, p^m)$ is convergent to zero when $N \rightarrow \infty$ that proves Lemma. $\square$

Summarize the results of four Lemmas 1, 2, 3 and 4 on convergence of the numerical series $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$.

Theorem 1 *Let a function $f(k) \in \mathbb{B}\{G_1(N); G_2(N)\}$ be given with a_i and b_i satisfying (1.6) and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K}p^{\gamma r}$ for all $r \geq r_*$. Then*

$$R_\infty(f, p^m) = \sum_{r=0}^{\infty} \frac{L_r(f; p^m)}{p^{a_1 r}}. \quad (2.39)$$

Proof According to Lemmas 1 and 2 if there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K}p^{\gamma r}$ for all $r \geq r_*$ then $\mathcal{R}_1(f; N, p^m)$ is convergent to $\sum_{r=0}^{\infty} L_r(f; p^m) p^{-a_1 r}$ in the whole range (1.6) of varying parameters a_1, b_1 and a_2, b_2 . On the other hand, according to Lemmas 3 and 4 by the same sufficient conditions the numerical series $\mathcal{R}_2(f; N, p^m)$ is convergent to zero in the same range (1.6) of varying parameters. Then, in accordance with (2.4) we arrive at (2.39). $\square$

There are a few questions on convergence of $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ which have been left open beyond the scope of Theorem 1. First, this is a problem of necessary convergence conditions which need further discussion. Another question arises in view of convergence to zero of $\mathcal{R}_2(f; N, p^m)$: it can happen that for some $f(k)$ both renormalization functions $\mathcal{R}_1(f; N, p^m)$ and $\mathcal{R}_2(f; N, p^m)$ are convergent to the non zero values while $L_r(f; p^m)$ is satisfying less strong conditions than those given in Theorem 1. Thus, the following question has been left still open,

Question 1 *Does the error term contribute to the renormalization function $R_\infty(f; p^m)$ and what multiplicative arithmetic functions $f(k)$ can provide an affirmative answer ?*

2.3 RATIONAL REPRESENTATION OF RENORMALIZATION FUNCTION

In sections 2.1 and 2.2 we have found the requirements which suffice to make $\mathcal{R}_1(f; N, p^m)$ convergent and $\mathcal{R}_2(f; N, p^m)$ vanishing. These conditions are presented through the characteristic functions $L_r(f; p^m)$ given recursively in (1.14). Their straightforward formulas (1.15) look cumbersome and lead in particular cases to rather complicate expressions, e.g., (1.17). This is why in this section we give another representation for $R_\infty(f; p^m)$ avoiding the use of $L_r(f; p^m)$.

Substituting (1.14) into (2.39) we get an infinite series for $R_\infty(f; p^m) \equiv R_\infty$,

$$R_\infty = f(p^m) + \frac{f(p^{m+1}) - L_0(f; p^m) f(p)}{p^{a_1}} + \frac{f(p^{m+2}) - L_1(f; p^m) f(p) - L_0(f; p^m) f(p^2)}{p^{2a_1}} + \frac{f(p^{m+3}) - L_2(f; p^m) f(p) - L_1(f; p^m) f(p^2) - L_0(f; p^m) f(p^3)}{p^{3a_1}} + \dots$$

Recasting the terms in the last expression we obtain

$$R_\infty(f; p^m) = \sum_{r=0}^{\infty} \frac{f(p^{m+r})}{p^{a_1 r}} - \frac{f(p)}{p^{a_1}} \sum_{r=0}^{\infty} \frac{L_r(f; p^m)}{p^{a_1 r}} - \frac{f(p^2)}{p^{2a_1}} \sum_{r=0}^{\infty} \frac{L_r(f; p^m)}{p^{a_1 r}} - \dots \quad (2.40)$$

Introduce two numerical series $\mathbf{U}(f; p^m)$ and $\mathbf{V}(f; p)$

$$\mathbf{U}(f; p^m) = \sum_{r=0}^{\infty} \frac{f(p^{r+m})}{p^{a_1 r}}, \quad \mathbf{V}(f; p) = \sum_{r=1}^{\infty} \frac{f(p^r)}{p^{a_1 r}}, \quad (2.41)$$

and assume that they are convergent. Then, by comparison of (2.40) and (2.39) we get

$$R_\infty(f; p^m) = \frac{\mathbf{U}(f; p^m)}{1 + \mathbf{V}(f; p)}. \quad (2.42)$$

if the denominator in (2.42) does not vanish. For short it can be written as follows, $\mathbf{V}(f; p) + 1 = \sum_{r=0}^{\infty} f(p^r) p^{-a_1 r}$. However, we prefer to stay with (2.42), otherwise one can make an error in calculations, e.g., $\phi(p^r) = p^{r-1}(p-1)$, $r \geq 1$, and $\phi(p^0) = 1$, but $\phi(p^0) \neq p^{-1}(p-1)$.

Formula (2.42) gives a rational representation of the renormalization function $R_\infty(f; p^m)$ which is free of intermediate calculations of $L_r(f; p^m)$. What can be said about convergence of $R_\infty(f; p^m)$ in terms of $f(p^r)$?

Theorem 2 *Let a function $f(k) \in \mathbb{B}\{G_1(N); G_2(N)\}$ be given and let there exist two numbers $\mathcal{K}_1 > 0$ and $\gamma_1 < a_1$ and an integer $r_* \geq 0$ such that $|f(p^r)| \leq \mathcal{K}_1 p^{\gamma_1 r}$ for all $r \geq r_*$. Then $R_\infty(f; p^m)$ is convergent in accordance with (2.42) if $\mathbf{V}(f; p) + 1 \neq 0$.*

Proof The constraints $|f(p^r)| \leq \mathcal{K}_1 p^{\gamma_1 r}$, $\mathcal{K}_1 > 0$ and $\gamma_1 < a_1$, for all $r \geq r_*$ implies the convergence of $\mathbf{U}(f; p^m)$ and $\mathbf{V}(f; p)$ that follows by their absolute convergence,

$$|\mathbf{U}(f; p^m)| \leq \mathcal{K}_1 p^{\gamma_1 m} \sum_{r=0}^{\infty} p^{(\gamma_1 - a_1)r} = \frac{\mathcal{K}_1 p^{\gamma_1 m}}{1 - p^{\gamma_1 - a_1}}, \quad |\mathbf{V}(f; p)| \leq \mathcal{K}_1 \sum_{r=0}^{\infty} p^{(\gamma_1 - a_1)r} = \frac{\mathcal{K}_1}{1 - p^{\gamma_1 - a_1}}.$$

Thus, if the denominator in (2.42) does not vanish, $\mathbf{V}(f; p) + 1 \neq 0$, then $R_\infty(f; p^m)$ is convergent in accordance with (2.42). $\square$

In Table 3 we present different multiplicative functions and their corresponding parameters a_1 , γ_1 , $\mathcal{K}_1$ and r_* . All functions satisfy the constraints of Theorem 2 for $p \geq 2$.

Table 3. Multiplicative functions and their corresponding parameters a_1 , γ_1 , $\mathcal{K}_1$ and r_* .

$f(k)$	$\phi(k)$	$1/\phi(k)$	$\psi(k)$	$1/\psi(k)$	$J_n(k)$	$\mu^2(k)$	$2^{\omega(k)}$	$3^{\omega(k)}$
a_1	2	0	2	0	$n+1$	1	1	1
γ_1	1	-1	1	-1	n	0	0	0
$\mathcal{K}_1$	$(p-1)/p$	$p/(p-1)$	$(p+1)/p$	$p/(p+1)$	$(p^n-1)/p^n$	1	2	3
r_*	1	1	1	1	1	0	0	0

Regarding the convergence of $\mathbf{U}(f; p^m)$ and $\mathbf{V}(f; p)$ defined in (2.41) we present an example which shows that the renormalization function $R_\infty(f; p^m)$ can exist even when both $\mathbf{U}(f; p^m)$ and $\mathbf{V}(f; p)$ are divergent. Consider $f(k) = 2^{\Omega(k)}$, $2^{\Omega(p^r)} = 2^r$ and get

$$\begin{cases} a_1 = \gamma_1 = 1 \\ \mathcal{K}_1 = 1 \end{cases}, \begin{cases} L_0(2^{\Omega(k)}; p^m) = 2^m \\ L_r(2^{\Omega(k)}; p^m) = 0, r \geq 1 \end{cases}, \begin{cases} \mathbf{U}(f; 2^m) \stackrel{N \rightarrow \infty}{\simeq} \infty \\ \mathbf{V}(f; 2) \stackrel{N \rightarrow \infty}{\simeq} \infty \end{cases}, R_\infty(2^{\Omega(k)}, 2^m) = 2^m.$$

Here the conditions of Theorem 1 are satisfied for all $r \geq 1$ and only the 1st term is left nonzero in series (2.39). Both $\mathbf{U}(f; 2^m)$ and $\mathbf{V}(f; 2)$ are divergent, e.g., $\mathbf{V}(f; 2) = \sum_{r=1}^{\infty} 1$, and therefore Theorem 2 cannot be applied. In other words, Theorem 1 has much wider area of application than Theorem 2. In the following sections we make use of both Theorems.

3 MULTIPLICATIVITY OF RENORMALIZATION FUNCTION WITH COMPLEX SCALING

In this section we study the renormalization of summatory function when the summation variable is scaled by a product $\prod_{i=1}^n p_i^{m_i}$ with n distinct primes $p_i \geq 2$. Consider $F\{f; N, \prod_{i=1}^n p_i^{m_i}\}$ and find its governing functional equation (3.2).

First, write the relationship between two summatory functions with two different summands, $f(k_1 p_n^{m_n})$ and $f(k_1)$, where $k_1 = k \prod_{i=1}^{n-1} p_i^{m_i}$. It is similar to that given in (1.13) and follows from the latter by replacing $k \rightarrow k_1$, i.e.,

$$F\left\{f; N, \prod_{i=1}^n p_i^{m_i}\right\} = \sum_{r_n=0}^{p_n^{r_n} \leq N} L_{r_n}(f; p_n^{m_n}) F\left\{f; \left\lfloor \frac{N}{p_n^{r_n}} \right\rfloor, \prod_{i=1}^{n-1} p_i^{m_i}\right\}. \quad (3.1)$$

Next, repeat this procedure to reduce the scale by $p_{n-1}^{r_{n-1}}$ for summatory function appeared in the r.h.s. of (3.1) and substitute it again into (3.1),

$$F\left\{f; N, \prod_{i=1}^n p_i^{m_i}\right\} = \sum_{r_{n-1}, r_n=0}^{p_{n-1}^{r_{n-1}} p_n^{r_n} \leq N} L_{r_n}(f; p_n^{m_n}) L_{r_{n-1}}(f; p_{n-1}^{m_{n-1}}) F\left\{f; \left\lfloor \frac{N}{p_{n-1}^{r_{n-1}} p_n^{r_n}} \right\rfloor, \prod_{i=1}^{n-2} p_i^{m_i}\right\}.$$

Continue to reduce the scales in a consecutive way for the next summatories and get finally,

$$F\left\{f; N, \prod_{i=1}^n p_i^{m_i}\right\} = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \left(\prod_{i=1}^n L_{r_i}(f; p_i^{m_i}) \right) F\left\{f; \left\lfloor \frac{N}{\prod_{i=1}^n p_i^{r_i}} \right\rfloor, 1\right\}. \quad (3.2)$$

Define new renormalization functions,

$$R\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) = \frac{F\{f; N, \prod_{i=1}^n p_i^{m_i}\}}{F\{f; N, 1\}}, \quad R_\infty\left(f; \prod_{i=1}^n p_i^{m_i}\right) := \lim_{N \rightarrow \infty} R\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) \quad (3.3)$$

and find their representations through the characteristic functions $L_r(f; p_i^{m_i})$, $i = 1, \dots, n$, and degrees a_1, b_1 and a_2, b_2 of leading asymptotics $G_1(N)$ and $G_2(N)$, respectively. Following an approach developed in section 2, represent $R(f; N, \prod_{i=1}^n p_i^{m_i})$ as a sum

$$R\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) = \mathcal{R}_1\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) + \mathcal{R}_2\left(f; N, \prod_{i=1}^n p_i^{m_i}\right), \quad (3.4)$$

where $\mathcal{R}_j(f; N, \prod_{i=1}^n p_i^{m_i})$, $j = 1, 2$, are analogous to those given in (2.2) and (2.3). Substitute there $G_1(N)$ and $G_2(N)$ given in (1.6), and obtain formulas analogous to those given in (2.5) and (2.6), (2.7). Here they are

$$\mathcal{R}_1\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \prod_{i=1}^n \left(\frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) \left(1 - \frac{1}{\log_c N} \sum_{i=1}^n r_i \log_c p_i \right)^{b_1}, \quad (3.5)$$

where a base c is chosen in such a way that $2 \leq c < \min\{p_1, \dots, p_n\}$, so that the upper summation bound $\prod_{i=1}^n p_i^{r_i} \leq N$ is correspondent to inequality, $\sum_{j=1}^n r_i \log_c p_i \leq \log_c N$. Regarding $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$, we have an upper bound

$$\left| \mathcal{R}_2\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) \right| \leq \frac{\mathcal{C}}{N^{a_2} (\log_c N)^{b_2}} \mathcal{R}_3\left(f; N, \prod_{i=1}^n p_i^{m_i}\right), \quad \text{where} \quad (3.6)$$

$$\mathcal{R}_3\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \prod_{i=1}^n \left(\frac{|L_{r_i}(f; p_i^{m_i})|}{p_i^{r_i(a_1 - a_2)}} \right) \left(1 - \frac{1}{\log_c N} \sum_{i=1}^n r_i \log_c p_i \right)^{b_1 - b_2}. \quad (3.7)$$

Keeping in mind the SQ theorem and its usage in sections 2.1 and 2.2 we assume throughout this section $b_1, b_2 \in \mathbb{Z}$. Extension on non integers b_1 and b_2 is trivial and can be done following those given in Lemmas 1 and 2, and therefore will be skipped. In next sections we prove several statements on $\mathcal{R}_1(f; N, \prod_{i=1}^n p_i^{m_i})$ and $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$ which are similar to Lemmas 1, 2, 3 and 4 in section 2. In this conjunction, it is important to use the same sufficient conditions which were used in these Lemmas.

3.1 CONVERGENCE OF $\mathcal{R}_1(f; N, \prod_{i=1}^n p_i^{m_i})$

Lemma 5 *Let a function $f(k) \in \mathbb{B}\{N^{a_1} (\log N)^{b_1}; G_2(N)\}$, $a_1 \geq 0$, $b_1 \in \mathbb{Z}_+ \cup \{0\}$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$. Then*

$$\lim_{N \rightarrow \infty} \mathcal{R}_1\left(f; N, \prod_{i=1}^n p_i^{m_i}\right) = \prod_{i=1}^n \left(\sum_{r_i=0}^{\infty} \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right). \quad (3.8)$$

Proof Exponentiating the binomial in (3.5) we obtain

$$\mathcal{R}_1 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) + \sum_{k=1}^{b_1} (-1)^k \binom{b_1}{k} \mathcal{R}_6 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right), \quad (3.9)$$

where

$$\mathcal{R}_6 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) = \frac{1}{(\log_c N)^k} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) \left(\sum_{i=1}^n r_i \log_c p_i \right)^k.$$

Find an estimate for $\mathcal{R}_6(f; N, \prod_{i=1}^n p_i^{m_i}, k)$,

$$\left| \mathcal{R}_6 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) \right| \leq \mathcal{K}^n \sum_{\substack{k_1, \dots, k_n=0 \\ k_1 + \dots + k_n = k}}^k \binom{k}{k_1, \dots, k_n} \prod_{i=1}^n \left(\frac{\log_c p_i}{\log_c N} \right)^{k_i} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \left(\prod_{i=1}^n \frac{r_i^{k_i}}{p_i^{\epsilon r_i}} \right),$$

where $\epsilon = a_1 - \gamma > 0$. One more inequality reads

$$\sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \left(\prod_{i=1}^n \frac{r_i^{k_i}}{p_i^{\epsilon r_i}} \right) \leq \prod_{i=1}^n \left(\sum_{r_i=0}^{p_i^{r_i} \leq N} \frac{r_i^{k_i}}{p_i^{\epsilon r_i}} \right) = \prod_{i=1}^n T(p_i, k_i, \epsilon, \log_{p_i} N), \quad (3.10)$$

where $T(p, k, \epsilon, M)$ is defined in (2.13). Combining the two last inequalities together we get

$$\left| \mathcal{R}_6 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) \right| \leq \mathcal{K}^n \sum_{\substack{k_1, \dots, k_n=0 \\ k_1 + \dots + k_n = k}}^k \binom{k}{k_1, \dots, k_n} \prod_{i=1}^n \frac{T(p_i, k_i, \epsilon, \log_{p_i} N)}{(\log_{p_i} N)^{k_i}}. \quad (3.11)$$

Inserting the asymptotics (2.15) of $T(p_i, k_i, \epsilon, \log_{p_i} N)$ into (3.11) we arrive at

$$\left| \mathcal{R}_6 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) \right| \leq \mathcal{K}^n \sum_{\substack{k_1, \dots, k_n=0 \\ k_1 + \dots + k_n = k}}^k \binom{k}{k_1, \dots, k_n} \prod_{i=1}^n \left[\frac{\text{Li}_{-k_i}(p_i^{-\epsilon})}{(\log_{p_i} N)^{k_i}} - \frac{N^{-\epsilon}}{p_i^{\epsilon} - 1} \right]. \quad (3.12)$$

Repeating the concluding remarks in proof of Lemma 1 on asymptotics of the polylogarithm function $\text{Li}_s(z)$ we conclude that $\mathcal{R}_6(f; N, \prod_{i=1}^n p_i^{m_i}, k)$ is convergent to zero when $N \rightarrow \infty$. Then, keeping in mind the representation (3.9) for $\mathcal{R}_1(f; N, \prod_{i=1}^n p_i^{m_i})$ and running the upper bound of summation to infinity we conclude that the limit (3.8) holds. $\square$

Lemma 6 *Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; G_2(N) \right\}$, $a_1 > 0$, $b_1 \in \mathbb{Z}_-$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$. Then (3.8) holds.*

Proof Here we follow the keyline in the proof of Lemma 2 and, according to (1.18) in Remark 2, start with representation of $\mathcal{R}_1(f; N, \prod_{i=1}^n p_i^{m_i})$ avoiding its divergence at $\prod_{i=1}^n p_i^{m_i} = N$,

$$\mathcal{R}_1 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N-1} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) \left(1 - \frac{1}{\log_c N} \sum_{i=1}^n r_i \log_c p_i \right)^{-|b_1|}. \quad (3.13)$$

Making use of identity (2.19) we obtain,

$$\mathcal{R}_1 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) + \sum_{k=1}^{|b_1|} \mathcal{R}_7 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right), \quad (3.14)$$

where $\tilde{r} = \sum_{i=1}^n r_i \log_c p_i$ and

$$\mathcal{R}_7 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) = \frac{1}{\log_c N} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right) \tilde{r} \left(1 - \frac{\tilde{r}}{\log_c N} \right)^{-k}. \quad (3.15)$$

Making use of inequality (2.22) and constraint imposed on $L_r(f; p^m)$

$$\left(1 - \frac{\tilde{r}}{\log_c N} \right)^{-k} \leq (1 + \tilde{r})^k, \quad |L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r},$$

exponentiate the binomial $(1 + \tilde{r})^k$ in (3.15) and obtain

$$\left| \mathcal{R}_7 \left(f; N, \prod_{i=1}^n p_i^{m_i}, k \right) \right| \leq \frac{\mathcal{K}^n}{\log_c N} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \frac{\tilde{r} (1 + \tilde{r})^k}{\prod_{i=1}^n p_i^{\epsilon r_i}} = \frac{\mathcal{K}^n}{\log_c N} \sum_{j=0}^k \binom{k}{j} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \frac{\tilde{r}^{j+1}}{p_1^{\epsilon r_1} p_2^{\epsilon r_2}}.$$

where $\epsilon = a_1 - \gamma$. Exponentiating a binomial $(\sum_{i=1}^n r_i \log_c p_i)^{j+1}$ in the last expression we get

$$|\mathcal{R}_7(f; N, p_1^{m_1} p_2^{m_2}, k)| \leq \frac{\mathcal{K}^n}{\log_c N} \sum_{j=0}^k \binom{k}{j} \sum_{\substack{j_1, \dots, j_n=0 \\ j_1 + \dots + j_n = j+1}}^{j+1} \binom{j+1}{j_1, \dots, j_n} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n \frac{r_i^{j_i}}{p_i^{\epsilon r_i}} (\log_c p_i)^{j_i},$$

Asymptotic behavior in N of the last expression is completely determined by its inner sum in r_i with respect to its prefactor $(\log_c N)^{-1}$. This behavior can be calculated following corresponding part (3.10) of the proof in Lemma 1,

$$\sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n \frac{r_i^{j_i}}{p_i^{\epsilon r_i}} (\log_c p_i)^{j_i} \stackrel{N \rightarrow \infty}{\leq} \prod_{i=1}^n \left[\text{Li}_{-j_i}(p_i^{-\epsilon}) - \frac{N^{-\epsilon} (\log_{p_i} N)^{j_i}}{1 - p_i^{-\epsilon}} \right] (\log_c p_i)^{j_i}. \quad (3.16)$$

Keeping in mind the prefactor $(\log_c N)^{-1}$ and the last asymptotics (3.16), the upper bound for $\mathcal{R}_7(f; N, \prod_{i=1}^n p_i^{m_i}, k)$ can be done infinitely small, i.e., it is convergent to zero when $N \rightarrow \infty$.

Making use of representation (3.14) for $\mathcal{R}_1(f; N, p_1^{m_1} p_2^{m_2})$ and running the upper bound of summation to infinity we conclude that the limit (3.8) holds. $\square$

3.2 CONVERGENCE OF $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$

Lemma 7 *Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$, $b_1 \geq b_2$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$, then*

$$\lim_{N \rightarrow \infty} \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = 0. \quad (3.17)$$

Proof Denote $b_1 - b_2 = e \in \mathbb{Z}_+ \cup \{0\}$ and make use of an inequality, $(1 - x)^e \leq 1$ when $0 \leq x \leq 1$, $e \geq 0$. Substituting constraints on $L_r(f; p^m)$ into (3.6) we obtain,

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \right| \leq \frac{\mathcal{CK}^n \mathcal{I}_n(\nu)}{N^{a_2} (\log_p N)^{b_2}}, \quad \text{where} \quad \mathcal{I}_n(\nu) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \prod_{i=1}^n p_i^{\nu r_i}. \quad (3.18)$$

and $\nu = a_2 - a_1 + \gamma$. Focus on the sum $\mathcal{I}_n(\nu)$ and estimate it in 3 cases, $\nu = 0$, $\nu > 0$ and $\nu < 0$.

Let $\nu = 0$, i.e., $a_2 = a_1 - \gamma$, then $\mathcal{I}_n(0)$ accounts for a number of integral points (vertices with integer coordinates) in the n -dim simplex, or corner of the n -dim cube, defined as follows,

$$\Delta_n =: \left\{ r_1, \dots, r_n \in \mathbb{Z}_+ \cup \{0\} \mid \sum_{i=1}^n r_i \log_c p_i \leq \log_c N \right\}.$$

The simplex Δ_n has one orthogonal corner and sizes of edges $\log_c N / \log_c p_i$ along the i th axis. The number $\mathcal{I}_n(0)$ is described by the Ehrhart polynomial [8] and, when $N \rightarrow \infty$, it has a leading term coinciding with simplex' volume.

$$\mathcal{I}_n(0) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{r_i} \leq N} \simeq \frac{(\log_c N)^n}{n! \prod_{i=1}^n \log_c p_i} \quad (3.19)$$

Substituting (3.19) into (3.18) we get

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{r_i} \right) \right| \leq \frac{\mathcal{CK}^n}{n! \prod_{i=1}^n \log_c p_i} \frac{N^{\gamma - a_1}}{(\log_c N)^{b_2 - n}}. \quad (3.20)$$

By (3.20) and inequality $\gamma < a_1$ we conclude that $\mathcal{R}_2(f; N, p_1^{m_1} p_2^{m_2})$ is convergent to zero when $N \rightarrow \infty$ irrespectively to the value of b_2 .

Consider the case $\nu > 0$ and estimate $\mathcal{I}_n(\nu)$ and $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{r_i})$,

$$\mathcal{I}_n(\nu) \leq N^\nu \mathcal{I}_n(0), \quad \rightarrow \quad \left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{r_i} \right) \right| \leq \frac{\mathcal{CK}^n}{n! \prod_{i=1}^n \log_c p_i} \frac{N^{\gamma - a_1}}{(\log_c N)^{b_2 - n}}. \quad (3.21)$$

By (3.21) the term $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{r_i})$ is also convergent to zero when $N \rightarrow \infty$.

Finally, consider the case $\nu < 0$. According to (3.18) we have $\mathcal{I}_n(\nu) < \mathcal{I}_n(0)$ and therefore

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{r_i} \right) \right| \leq \frac{\mathcal{CK}^n}{n! \prod_{i=1}^n \log_c p_i} \frac{N^{-a_2}}{(\log_c N)^{b_2 - n}}. \quad (3.22)$$

Thus, $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{r_i})$ is convergent to zero due to (3.22) and constraints (1.6) on degrees a_2 and b_2 . Summarizing (3.20), (3.21) and (3.22) we complete the proof of Lemma. $\square$

In the following Lemmas we consider two different cases, $a_2 > 0$ and $a_2 = 0$, separately.

Lemma 8 Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; N^{-a_2} (\log N)^{-b_2} \right\}$, $b_1 < b_2$ and $a_2 > 0$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$. Then (3.17) holds.

Proof In accordance with (1.18) in Remark 2, represent $\mathcal{R}_3(f; N, \prod_{i=1}^n p_i^{m_i})$ avoiding its divergence at $\prod_{i=1}^n p_i^{m_i} = N$,

$$\mathcal{R}_3 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \left(\prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{r_i(a_1-a_2)}} \right) \left(1 - \frac{\tilde{r}}{\log_c N} \right)^{-(b_2-b_1)}, \quad (3.23)$$

and make use of inequality (2.22) in the range $0 \leq \tilde{r} \leq \lfloor \log_p N \rfloor - 1$,

$$\left(1 - \frac{\tilde{r}}{\log_c N} \right)^{-(b_2-b_1)} \leq (\log_c N)^{b_2-b_1}. \quad (3.24)$$

Substituting (3.24) into (3.23) for $\mathcal{R}_3(f; N, \prod_{i=1}^n p_i^{m_i})$ we get

$$\mathcal{R}_3 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \leq (\log_c N)^{b_2-b_1} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n \frac{|L_{r_i}(f; p_i^{m_i})|}{p_i^{r_i(a_1-a_2)}}. \quad (3.25)$$

Apply to (3.25) the constraints on $L_r(f; p^m)$ and substitute the result into (3.6),

$$\mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \leq \frac{\mathcal{CK}^n}{N^{a_2} (\log_c N)^{b_1}} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n p_i^{\nu r_i}, \quad \nu = a_2 - a_1 + \gamma. \quad (3.26)$$

By comparison (3.26) with (3.18) from Lemma 7 we obtain according to (3.20), (3.21) and (3.22)

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \right| \leq \frac{\mathcal{CK}^n}{n! \prod_{i=1}^n \log_c p_i} \frac{N^{\gamma-a_1}}{(\log_c N)^{b_1-n}}, \quad \nu \geq 0, \quad (3.27)$$

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \right| \leq \frac{\mathcal{CK}^n}{n! \prod_{i=1}^n \log_c p_i} \frac{N^{-a_2}}{(\log_c N)^{b_1-n}}, \quad \nu < 0. \quad (3.28)$$

Thus, by (3.27) and inequality $\gamma < a_1$ a series $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$ is convergent to zero when $N \rightarrow \infty$ irrespectively to the value of b_1 . The same conclusion (3.17) on convergence of this series holds due to (3.28) when $a_2 > 0$. $\square$

Lemma 9 Let a function $f(k) \in \mathbb{B} \left\{ N^{a_1} (\log N)^{b_1}; (\log N)^{-b_2} \right\}$, $b_1 < b_2$, $a_2 = 0$, be given and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K} p^{\gamma r}$ for all $r \geq r_*$. Then (3.17) holds.

Proof This case has to be treated more precisely than that in Lemma 8. Rewrite (3.23)

$$\mathcal{R}_3 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) = \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n \frac{|L_{r_i}(f; p_i^{m_i})|}{p_i^{a_1 r_i}} \left(1 - \frac{\tilde{r}}{\log_c N} \right)^{-(b_2-b_1)}, \quad (3.29)$$

and compare it with expression (3.13) when $b_1 < 0$. A difference in degrees, $b_2 - b_1$ and b_1 , does not break the main result of Lemma 6 : only the 1st leading term in (3.14) is survived when $N \rightarrow \infty$. When we apply it to (3.29) and make use of constraint on $L_r(f; p^m)$ we get

$$\mathcal{R}_3 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \stackrel{N \rightarrow \infty}{\asymp} \sum_{r_1, \dots, r_n=0}^{\prod_{i=1}^n p_i^{m_i} \leq N-1} \prod_{i=1}^n \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \leq \mathcal{K}^n \prod_{i=1}^n \left(\sum_{r_i=0}^{\infty} p_i^{-\epsilon r_i} \right) = \frac{\mathcal{K}^n}{\prod_{i=1}^n (1 - p_i^{-\epsilon})},$$

where $\epsilon = a_1 - \gamma > 0$. Substituting the last estimate into (3.6) we obtain,

$$\left| \mathcal{R}_2 \left(f; N, \prod_{i=1}^n p_i^{m_i} \right) \right| \leq \frac{\mathcal{C}\mathcal{K}^n}{\prod_{i=1}^n (1 - p_i^{-\epsilon})} \cdot \frac{1}{(\log_p N)^{b_2}}. \quad (3.30)$$

Recall that by (1.6) the degrees of the error term satisfy the condition: if $a_2 = 0$ than $b_2 > 0$. Thus, by (3.30) the series $\mathcal{R}_2(f; N, \prod_{i=1}^n p_i^{m_i})$ converges to zero when $N \rightarrow \infty$ that proves Lemma. $\square$

We combine Lemmas 5, 6, 7, 8 and 9 on convergence of $\mathcal{R}_j(f; N, \prod_{i=1}^n p_i^{m_i})$, $j = 1, 2$, and according to (3.4) we arrive at the analogue of Theorem 1 in the case of a scaling by $\prod_{i=1}^n p_i^{m_i}$.

Theorem 3 *Let a function $f(k) \in \mathbb{B}\{G_1(N); G_2(N)\}$ be given with a_i and b_i satisfying (1.6) and let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(f; p^m)| \leq \mathcal{K}p^{\gamma r}$ for all $r \geq r_*$. Then*

$$R_\infty \left(f; \prod_{i=1}^n p_i^{m_i} \right) = \prod_{i=1}^n \left(\sum_{r_i=0}^{\infty} \frac{L_{r_i}(f; p_i^{m_i})}{p_i^{a_1 r_i}} \right). \quad (3.31)$$

Combining Theorems 1 and 3 we come to important consequence which manifests the multiplicative property of the renormalization function.

Corollary 1 *Under the conditions of Theorem 3 the following holds,*

$$R_\infty \left(f; \prod_{i=1}^n p_i^{m_i} \right) = \prod_{i=1}^n R_\infty(f; p_i^{m_i}). \quad (3.32)$$

3.3 ASYMPTOTICS OF SUMMATORY FUNCTIONS $\sum_{k_1, k_2 \leq N} f(k_1 k_2)$

In this section we calculate the summatory function $\Phi[f; N, 1] = \sum_{k_1, k_2 \leq N} f(k_1 k_2)$ and find its asymptotics by applying Corollary 1. According to definition of summatory function we get,

$$\Phi\{f; N, 1\} = \sum_{k \leq N} f(k) + \sum_{k \leq N} f(2k) + \dots = \sum_{k \leq N} \sum_{\substack{m_{i,j}=0 \\ n_j=1}} f \left(k \prod_{i=1}^{n_j} p_i^{m_{i,j}} \right) = \sum_{\substack{m_{i,j}=0 \\ n_j=1}} \sum_{k \leq N} f \left(k \prod_{i=1}^{n_j} p_i^{m_{i,j}} \right)$$

where indices $i, j, m_{i,j}$ and n_j account for all primes such that $\prod_{i=1}^{n_j} p_i^{m_{i,j}} \leq N$. Thus, according to definition of summatory function with scaled summation variable, we obtain,

$$\Phi\{f; N, 1\} = \sum_{\substack{m_{i,j}=0 \\ n_j=1}} F \left\{ f; N, \prod_{i=1}^{n_j} p_i^{m_{i,j}} \right\}, \quad \frac{\Phi\{f; N, 1\}}{F\{f; N, 1\}} = \sum_{\substack{m_{i,j}=0 \\ n_j=1}} R \left(f; N, \prod_{i=1}^{n_j} p_i^{m_{i,j}} \right), \quad (3.33)$$

where $\prod_{i=1}^{n_j} p_i^{m_{i,j}} \leq N$. Consider asymptotics (omitting the error terms) of three summatory functions when $N \rightarrow \infty$,

$$\begin{aligned} F\{f; N, 1\} &\stackrel{N \rightarrow \infty}{=} \mathcal{F}G_1(N), & \Phi\{f; N, 1\} &\stackrel{N \rightarrow \infty}{=} \mathcal{G}_1(f)\Gamma_1(N), \\ \sum_{\substack{m_{i,j}=0 \\ n_j=1}} R\left(f; N, \prod_{i=1}^{n_j} p_i^{m_{i,j}}\right) &\stackrel{N \rightarrow \infty}{=} \mathcal{G}_2(f)\Gamma_2(N). \end{aligned} \quad (3.34)$$

Combining (3.34) and the 2nd formula in (3.33) we obtain

$$\Gamma_1(N) = G_1(N) \cdot \Gamma_2(N), \quad \mathcal{G}_1(f) = \mathcal{F} \cdot \mathcal{G}_2(f). \quad (3.35)$$

Calculation of $\Gamma_2(N)$ and $\mathcal{G}_2(f)$ is a difficult numerical task. Consider a special case when $\mathcal{G}_2(f)$ may be given in a closed form, namely, when $\Gamma_2(N) = N^0$, i.e., $\Gamma_1(N) = G_1(N)$. Consider the 3rd asymptotics in (3.34) and, according to Corollary 1, find its limit when $N \rightarrow \infty$,

$$\begin{aligned} \mathcal{G}_2(f) &= \lim_{N \rightarrow \infty} \sum_{\substack{\prod_{i=1}^{n_j} p_i^{m_{i,j}} \leq N \\ m_{i,j}=0 \\ n_j=1}} R\left(f; N, \prod_{i=1}^{n_j} p_i^{m_{i,j}}\right) = \lim_{\substack{N \rightarrow \infty \\ n_j \rightarrow \infty}} \sum_{\substack{m_{i,j}=0 \\ n_j=1}}^{\log_{p_i} N} \prod_{i=1}^{n_j} R_\infty(f; p_i^{m_{i,j}}) \\ &= \lim_{n \rightarrow \infty} \prod_{i=1}^n \sum_{m=0}^{\infty} R_\infty(f; p_i^m) = \prod_{p \geq 2} \sum_{m=0}^{\infty} R_\infty(f; p^m), \end{aligned} \quad (3.36)$$

Another special case comes when $f(k)$ is a completely multiplicative arithmetic function. i.e., $f(k_1 k_2) = f(k_1) f(k_2)$. This leads to equalities: $\Gamma_2(N) \equiv \Gamma_1(N)$ and $\mathcal{G}_2(f) = \mathcal{F}$. We will illustrate this statement and (3.36) in section 4.2.

4 RENORMALIZATION OF DIRICHLET SERIES AND OTHERS SUMMATORY FUNCTIONS

In this section we extend the renormalization approach on summatory functions of more complex structure. They involve the summatory functions with summands given by $\prod_{i=1}^n f_i(k)$ and summation variable k scaled for every multiplicative function f_i by p^{m_i} , $m_i \neq m_j$. The case of the Dirichlet series is a special one when $n = 2$ and $f_2(k) = k^{-s}$. We study also the renormalization of summatory functions with summands given by $f(k^n)$.

4.1 RENORMALIZATION OF SUMMATORY FUNCTION $\sum_{k \leq N} \prod_{i=1}^n f_i(k p^{m_i})$

Start with summatory function $F\{\prod_{i=1}^n f_i; N, p^{\mathbf{m}}\} = \sum_{k \leq N} \prod_{i=1}^n f_i(k p^{m_i})$, where $p^{\mathbf{m}}$ denotes a tuple $\{p^{m_1}, \dots, p^{m_n}\}$, and make use of a standard notation $F\{\prod_{i=1}^n f_i; N, 1^{\mathbf{m}}\} = F\{\prod_{i=1}^n f_i; N, 1\}$.

Derive for F a functional equation following the approach developed in section 1.2 and start

$$\begin{aligned} F \left\{ \prod_{i=1}^n f_i; N, p^{\mathbf{m}} \right\} &= F(p^{\mathbf{m}}) \sum_{\substack{k=1 \\ p \nmid k}}^N \prod_{i=1}^n f_i(k) + \sum_{\substack{k=p \\ p \mid k}}^N \prod_{i=1}^n f_i(kp^{m_i}) \\ &= F(p^{\mathbf{m}}) \left(\sum_{k=1}^N \prod_{i=1}^n f_i(k) - \sum_{\substack{k=p \\ p \mid k}}^N \prod_{i=1}^n f_i(k) \right) + \sum_{l=1}^{N_1} \prod_{i=1}^n f_i(lp^{m_i+1}), \end{aligned}$$

where $F(p^{\mathbf{m}}) = \prod_{i=1}^n f_i(p^{m_i})$ and N_r was defined in section 1.2. Rewrite the last equality

$$F \left\{ \prod_{i=1}^n f_i; N, p^{\mathbf{m}} \right\} - F \left\{ \prod_{i=1}^n f_i; N_1, p^{\mathbf{m}+1} \right\} = F(p^{\mathbf{m}}) \left[F \left\{ \prod_{i=1}^n f_i; N, 1 \right\} - F \left\{ \prod_{i=1}^n f_i; N_1, p \right\} \right],$$

which is similar to (1.11). The corresponding counterpart for its general version (1.12) reads,

$$F \left\{ \prod_{i=1}^n f_i; N_r, p^{\mathbf{m}} \right\} - F \left\{ \prod_{i=1}^n f_i; N_{r+1}, p^{\mathbf{m}+1} \right\} = F(p^{\mathbf{m}}) \left[F \left\{ \prod_{i=1}^n f_i; N_r, 1 \right\} - F \left\{ \prod_{i=1}^n f_i; N_{r+1}, p \right\} \right]$$

Combining last equations of running index $0 \leq r \leq \lfloor \log_p N \rfloor$ together we arrive at the functional equation for summatory function,

$$F \left\{ \prod_{i=1}^n f_i; N, p^{\mathbf{m}} \right\} = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) F \left\{ \prod_{i=1}^n f_i; N_r, 1 \right\}, \quad (4.1)$$

where

$$L_r \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) = \prod_{i=1}^n f_i(p^{m_i+r}) - \sum_{j=0}^{r-1} L_j \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) \prod_{i=1}^n f_i(p^{r-j}), \quad (4.2)$$

Formulas for the first $L_r(\prod_{i=1}^n f_i; p^{\mathbf{m}})$ read

$$\begin{aligned} L_0 \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) &= \prod_{i=1}^n f_i(p^{m_i}), \quad L_1 \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) = \prod_{i=1}^n f_i(p^{m_i+1}) - \prod_{i=1}^n f_i(p^{m_i}) \prod_{i=1}^n f_i(p), \\ L_2 \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) &= \prod_{i=1}^n f_i(p^{m_i+2}) - \prod_{i=1}^n f_i(p^{m_i}) \prod_{i=1}^n f_i(p^2) - \prod_{i=1}^n f_i(p^{m_i+1}) \prod_{i=1}^n f_i(p) + \\ &\quad \prod_{i=1}^n f_i(p^{m_i}) \prod_{i=1}^n f_i^2(p), \quad \text{etc} \end{aligned} \quad (4.3)$$

such that for $p = 1$ or $m = 0$ we have, $L_0(\prod_{i=1}^n f_i; 1^{\mathbf{m}}) = 1$ and $L_r(\prod_{i=1}^n f_i; 1^{\mathbf{m}}) = 0$, $r \geq 1$. Find analogues to formulas (1.16) when $f_i(p^m) = A_{f_i} p^{m-1}$, $m \geq 1$, and A_{f_i} denotes the real constant. By (4.2) or (4.3) such formulas for $L_r(\prod_{i=1}^n f_i; p^{\mathbf{m}})$ can be calculated by induction,

$$\begin{aligned} L_r \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) &= p^{m_1+\dots+m_n-n} \left(p^n - \prod_{i=1}^n A_{f_i} \right)^r \prod_{i=1}^n A_{f_i}, \\ L_r \left(\prod_{i=1}^n \frac{1}{f_i}; p^{\mathbf{m}} \right) &= p^{n-m_1-\dots-m_n} \left(p^{-n} - \prod_{i=1}^n A_{f_i}^{-1} \right)^r \prod_{i=1}^n A_{f_i}^{-1}. \end{aligned}$$

E.g., in the case of the Euler $f_1 = \phi(k)$ and Dedekind $f_2 = \psi(k)$ totient functions we have

$$L_r(\phi\psi; p^{\mathbf{m}}) = (p^2 - 1)p^{m_1+m_2-2}, \quad L_r\left(\frac{1}{\phi\psi}; p^{\mathbf{m}}\right) = \frac{(-1)^r p^{2(1-r)-m_1-m_2}}{(p^2 - 1)^{r+1}}.$$

Define new renormalization functions,

$$R\left(\prod_{i=1}^n f_i; N, p^{\mathbf{m}}\right) = \frac{F\{\prod_{i=1}^n f_i; N, p^{\mathbf{m}}\}}{F\{\prod_{i=1}^n f_i; N, 1\}}, \quad R_{\infty}\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) = \lim_{N \rightarrow \infty} R\left(\prod_{i=1}^n f_i; N, p^{\mathbf{m}}\right). \quad (4.4)$$

By comparison formulas (4.1), (4.2), (4.3) with (1.13), (1.14), (1.15), respectively, and definition (4.4) with (1.3) we conclude that all results on renormalization of summatory function $F\{f; N, p^{\mathbf{m}}\}$ in section 2 can be reproduced for summatory function $F\{\prod_{i=1}^n f_i; N, 1^{\mathbf{m}}\}$ with a few necessary alterations. Below, in Theorem 4 we give (without proof) a sufficient condition for convergence of the asymptotics of renormalization function $R_{\infty}(\prod_{i=1}^n f_i; p^{\mathbf{m}})$. Its proof does not use new ideas and can be given following Lemmas 1, 2, 3 and 4 for renormalization function $R_{\infty}(f; p^{\mathbf{m}})$. For this reason we have skipped this proof here.

Theorem 4 *Let n multiplicative functions $f_i(k)$, be given such that $f_i(k) \in \mathbb{B}\{G_1(N); G_2(N)\}$ satisfying (1.6). Let there exist two numbers $\mathcal{K} > 0$ and $\gamma < a_1$ and an integer $r_* \geq 0$ such that $|L_r(\prod_{i=1}^n f_i; p^{\mathbf{m}})| \leq \mathcal{K}p^{\gamma r}$ for all $r \geq r_*$. Then*

$$R_{\infty}\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) = \sum_{r=0}^{\infty} L_r\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) p^{-a_1 r}. \quad (4.5)$$

To study $R_{\infty}(\prod_{i=1}^n f_i; p^{\mathbf{m}})$ in a way similar to the study of $R_{\infty}(f; p^{\mathbf{m}})$ in section 2 we find another representation for $R_{\infty}(\prod_{i=1}^n f_i; p^{\mathbf{m}})$ which is different from (4.5). Substitute (4.2) into (4.5) and get

$$\begin{aligned} R_{\infty}\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) &= \prod_{i=1}^n f_i(p^{m_i}) + \left[\prod_{i=1}^n f_i(p^{m_i+1}) - L_0\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) \prod_{i=1}^n f_i(p) \right] p^{-a_1} + \\ &\quad \left[\prod_{i=1}^n f_i(p^{m_i+2}) - L_1\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) \prod_{i=1}^n f_i(p) - L_0\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) \prod_{i=1}^n f_i(p^2) \right] p^{-2a_1} + \dots \end{aligned}$$

Recasting the terms in the last expression we obtain

$$R_{\infty}\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) = \sum_{r=0}^{\infty} p^{-a_1 r} \prod_{i=1}^n f_i(p^{m_i+r}) - \sum_{r=1}^{\infty} p^{-a_1 r} \prod_{i=1}^n f_i(p^r) \cdot \sum_{r=0}^{\infty} L_r\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) p^{-a_1 r}.$$

Thus, by comparison the last expression with formula (4.5) we get,

$$R_{\infty}\left(\prod_{i=1}^n f_i; p^{\mathbf{m}}\right) = \frac{\mathbf{U}(\prod_{i=1}^n f_i; p^{\mathbf{m}})}{1 + \mathbf{V}(\prod_{i=1}^n f_i; p)}, \quad (4.6)$$

where two numerical series

$$\mathbf{U} \left(\prod_{i=1}^n f_i; p^{\mathbf{m}} \right) = \sum_{r=0}^{\infty} p^{-a_1 r} \prod_{i=1}^n f_i(p^{m_i+r}) , \quad \mathbf{V} \left(\prod_{i=1}^n f_i; p \right) = \sum_{r=1}^{\infty} p^{-a_1 r} \prod_{i=1}^n f_i(p^r) \quad (4.7)$$

are assumed to be convergent and a denominator in (4.6) does not vanish. Formula (4.6) gives a rational representation of the renormalization function $R_{\infty}(\prod_{i=1}^n f_i; p^{\mathbf{m}})$ which is free of intermediate calculations of $L_r(\prod_{i=1}^n f_i; p^{\mathbf{m}})$.

4.2 RENORMALIZATION OF THE DIRICHLET SERIES $\sum_{k=1}^{\infty} f(kp^m) k^{-s}$

The Dirichlet series $D(f; p^m, s) = \sum_{k=1}^{\infty} f(kp^m) k^{-s}$ with a scaled summation variable is a special case of summatory function $F\{f_1 f_2; N, p^{\mathbf{m}}\}$, discussed in section 4.1, when $f_1 = f(k)$, $f_2 = k^{-s}$, $m_1 = m$, $m_2 = 0$ and $f_1 f_2 \in \mathbb{B}\{N^0; G_2(N)\}$, i.e., $a_1 = 0$.

According to Theorem 4 and formula (4.6) if there exist two numbers $\mathcal{K}_1 > 0$ and $\gamma_1 < 0$ and an integer $r_* \geq 0$ such that $|f(p^r)| \leq \mathcal{K}_1 p^{(s+\gamma_1)r}$ for all $r \geq r_*$ then

$$\mathfrak{D}(f; p^m, s) = \frac{D(f; p^m, s)}{D(f, s)} = \sum_{r=0}^{\infty} \frac{f(p^{r+m})}{p^{sr}} \left(1 + \sum_{r=1}^{\infty} \frac{f(p^r)}{p^{sr}} \right)^{-1}. \quad (4.8)$$

where $D(f, s) = \sum_{k=1}^{\infty} f(k) k^{-s}$ is a standard Dirichlet series for arithmetic function $f(k)$. Note that according to definition (4.4) of renormalization function $R_{\infty}(f \cdot k^{-s}; p^m)$ for the Dirichlet series the following equality holds, $\mathfrak{D}(f; p^m, s) = p^{sm} R_{\infty}(f \cdot k^{-s}; p^m)$.

We present four examples with the Dirichlet series $D(f; p^m, s)$ for the Möbius $\mu(k)$, Liouville $\lambda(k)$, Euler $\phi(k)$ and divisor $\sigma_n(k)$ functions. Their standard Dirichlet series $D(f, s)$ converge to the values given in Table 2.

- Consider two Dirichlet series for the μ -function, $D(\mu, s) = 1/\zeta(s)$ and $D(\mu^2, s) = \zeta(s)/\zeta(2s)$, $s > 1$, and calculate their scaled versions. Since $\mu(p^m k) \equiv 0$, $m \geq 2$, we consider here only a case $m = 1$ and have $\mu^q(p^{r+1}) = (-1)^q \delta_{0,r}$, $q = 1, 2$. Then

$$\mathfrak{D}(\mu; p, s) = -\frac{p^s}{p^s - 1}, \quad \mathfrak{D}(\mu^2; p, s) = \frac{p^s}{p^s + 1}. \quad (4.9)$$

Calculate the Dirichlet series $\Delta(\mu^q, s) = \sum_{k_1, k_2=1}^{\infty} (k_1 k_2)^{-s} \mu^q(k_1 k_2)$, $q = 1, 2$ according to (3.36)

$$\Delta(\mu, s) = \frac{1}{\zeta(s)} \mathcal{G}_2\left(\frac{\mu}{k^s}\right), \quad \mathcal{G}_2\left(\frac{\mu}{k^s}\right) = \prod_{p \geq 2} \left(1 + R_{\infty}\left(\frac{\mu}{k^s}; p\right)\right) = \prod_{p \geq 2} \left(1 - \frac{1}{p^s - 1}\right) < \frac{1}{\zeta(s)}. \quad (4.10)$$

$$\Delta(\mu^2, s) = \frac{\zeta(s)}{\zeta(2s)} \mathcal{G}_2\left(\frac{\mu^2}{k^s}\right), \quad \mathcal{G}_2\left(\frac{\mu^2}{k^s}\right) = \prod_{p \geq 2} \left(1 + R_{\infty}\left(\frac{\mu^2}{k^s}; p\right)\right) = \prod_{p \geq 2} \left(1 + \frac{1}{p^s + 1}\right) < \frac{\zeta(s)}{\zeta(2s)}$$

that results in inequalities, $\Delta(\mu, s) < D^2(\mu, s)$ and $\Delta(\mu^2, s) < D^2(\mu^2, s)$. A normalized product for $s = 2$ in (4.10) is known as the Feller-Tornier constant C_{FT} [16],

$$\frac{1}{\zeta(2)} \prod_{p \geq 2} \left(1 - \frac{1}{p^2 - 1}\right) = \prod_{p \geq 2} \left(1 - \frac{2}{p^2}\right) = C_{FT} = 0.32263 .$$

- Consider the Dirichlet series for the λ -function, $D(\lambda, s) = \zeta(2s)/\zeta(s)$, $s > 1$, and calculate its scaled version. Keeping in mind $\lambda(p^r) = (-1)^r$ we get $\mathfrak{D}(\lambda; p^m, s) = (-1)^m$. Calculate the Dirichlet series $\Delta(\lambda, s) = \sum_{k_1, k_2=1}^{\infty} (k_1 k_2)^{-s} \lambda(k_1 k_2)$ in accordance with (3.36)

$$\Delta(\lambda, s) = \frac{\zeta(2s)}{\zeta(s)} \mathcal{G}_2\left(\frac{\lambda}{k^s}\right), \quad \mathcal{G}_2\left(\frac{\lambda}{k^s}\right) = \prod_{p \geq 2} \sum_{m=0}^{\infty} (-1)^m p^{-sm} = \prod_{p \geq 2} \frac{1}{1 + p^{-s}} = \frac{\zeta(2s)}{\zeta(s)}, \quad (4.11)$$

i.e., $\Delta(\lambda, s) = D^2(\lambda, s)$ in accordance with the fact that $\lambda(k)$ is completely multiplicative.

- Consider the Dirichlet series for the ϕ -function, $D(\phi, s) = \zeta(s-1)/\zeta(s)$, $s > 2$, and calculate its scaled version. Keeping in mind $\phi(p^r) = (p-1)p^{r-1}$ we get

$$\mathfrak{D}(\phi; p^m, s) = \frac{(p-1)p^{m-1}}{1 - p^{-s}}, \quad m \geq 1. \quad (4.12)$$

- Consider two Dirichlet series for the σ_0 -function, $D(\sigma_0, s) = \zeta^2(s)$ and $D(\sigma_0^2, s) = \zeta^4(s)/\zeta(2s)$, $s \geq 1$, and calculate its scaled version. Keeping in mind $\sigma_0(p^r) = r+1$ we get

$$\begin{aligned} \mathfrak{D}(\sigma_0; p^m, s) &= (m+1)(1 - p^{-s}) + p^{-s}, \quad m \geq 0, \\ \mathfrak{D}(\sigma_0^2; p^m, s) &= \frac{[(m+1)(1 - p^{-s}) + p^{-s}]^2 + p^{-s}}{1 + p^{-s}}. \end{aligned} \quad (4.13)$$

We finish this section with relationship between characteristic functions for multiplicative arithmetic functions $f(k)$ and $f_1(k) = f(k) \cdot k^{-s}$

$$L_r(f \cdot k^{-s}; p^m) = L_r(f; p^m) p^{-(m+r)s}, \quad (4.14)$$

which follows by (1.14) and (1.15) if we substitute there the identity $f_1(p^r) = f(p^r) p^{-rs}$.

Relation (4.14) will be used in section 5.3 when calculating the renormalized Dirichlet series for the Ramanujan τ function.

4.3 RENORMALIZATION OF SUMMATORY FUNCTION $\sum_{k \leq N} f(k^n p^m)$

Consider the summatory function $F\{f, n; N, p^m\} = \sum_{k \leq N} f(k^n p^m)$ and derive its governing functional equation following the approach developed in section 1.2,

$$F\{f, n; N, p^m\} = f(p^m) \sum_{\substack{k=1 \\ p \nmid k}}^N f(k^n) + \sum_{\substack{k=p \\ p \mid k}}^N f(p^m k^n) = f(p^m) \left(\sum_{k=1}^N - \sum_{\substack{k=p \\ p \mid k}}^N \right) f(k^n) + \sum_{l=1}^{N_1} f(p^{m+n} l^n)$$

which can be rewritten as follows,

$$F\{f, n; N, p^m\} - F\{f, n; N_1, p^{m+n}\} = f(p^m) [F\{f, n; N, 1\} - F\{f, n; N_1, p^n\}] . \quad (4.15)$$

In general case ($r \geq 1$) an Eq. (4.15) has a form

$$F\{f, n; N_r, p^{m+rn}\} - F\{f, n; N_{r+1}, p^{m+(r+1)n}\} = f(p^{m+rn}) [F\{f, n; N_r, 1\} - F\{f, n; N_{r+1}, p^n\}]$$

Combining last equations of running index $0 \leq r \leq \lfloor \log_p N \rfloor$ together we arrive at the functional equation,

$$F\{f, n; N_r, p^m\} = \sum_{r=0}^{\lfloor \log_p N \rfloor} L_r(f, n; p^m) F\{f, n; N_r, 1\} , \quad \text{where} \quad (4.16)$$

$$L_r(f, n; p^m) = f(p^{m+rn}) - \sum_{j=0}^{r-1} L_j(f, n; p^m) f(p^{(r-j)n}) . \quad (4.17)$$

The straightforward calculations of $L_r(f; p^m)$ give

$$L_0(f, n; p^m) = f(p^m) , \quad L_1(f, n; p^m) = f(p^{m+n}) - f(p^m) f(p^n) , \quad (4.18)$$

$$L_2(f, n; p^m) = f(p^{m+2n}) - f(p^{m+n}) f(p^n) - f(p^m) [f(p^{2n}) - f^2(p^n)] ,$$

By $n = 1$ formulas (4.16), (4.17) and (4.18) are reduced to (1.13), (1.14) and (1.15).

Define new renormalization functions,

$$R(f, n; N, p^m) = \frac{F\{f, n; N, p^m\}}{F\{f, n; N, 1\}} , \quad R_\infty(f, n; p^m) := \lim_{N \rightarrow \infty} R(f, n; N, p^m) . \quad (4.19)$$

By comparison formulas (4.16), (4.17), (4.18) with (1.13), (1.14), (1.15), respectively, and definition (4.4) with (4.19) we conclude that all results on renormalization of summatory function $F\{f; N, p^m\}$ in section 2 can be reproduced for summatory function $F\{f, n; N, p^m\}$ with a few necessary alterations. Below we give (without proof) Theorem 5 on sufficient condition to converge of asymptotics of renormalization function $R_\infty(f, n; p^m)$. As in the case of Theorem 4 on renormalization function $R_\infty(f_1 f_2; p^m)$, here the proof of Theorem 5 does not use new ideas and can be given following Lemmas 1, 2, 3 and 4 for renormalization function $R_\infty(f; p^m)$. For this reason we skip it here.

Theorem 5 *Let a function $f(k^n) \in \mathbb{B}\{G_1(N); G_2(N)\}$ be given and let there exist two numbers $\mathcal{K}_1 > 0$ and $\gamma_1 < a_1$ and an integer $r_* \geq 0$ such that $|f(p^r)| \leq \mathcal{K}_1 p^{\gamma_1 r}$ for all $r \geq r_*$. Then*

$$R_\infty(f, n; p^m) = \sum_{r=0}^{\infty} \frac{L_r(f, n; p^m)}{p^{a_1 r}} . \quad (4.20)$$

Recasting the terms in the last expression we obtain

$$R_{\infty}(f, n; p^m) = \sum_{r=0}^{\infty} \frac{f(p^{m+nr})}{p^{a_1 r}} - \sum_{r=0}^{\infty} \frac{L_r(f, n; p^m)}{p^{a_1 r}} \left[\frac{f(p^n)}{p^{a_1}} + \frac{f(p^{2n})}{p^{2a_1}} + \frac{f(p^{3n})}{p^{3a_1}} + \dots \right].$$

Thus, by comparison the last expression with formula (4.5) we get,

$$R_{\infty}(f, n; p^m) = \frac{\mathbf{U}(f, n; p^m)}{1 + \mathbf{V}(f, n; p)}, \quad (4.21)$$

where two numerical series

$$\mathbf{U}(f, n; p^m) = \sum_{r=0}^{\infty} \frac{f(p^{m+nr})}{p^{a_1 r}}, \quad \mathbf{V}(f, n; p) = \sum_{r=1}^{\infty} \frac{f(p^{rn})}{p^{a_1 r}}, \quad (4.22)$$

are assumed to be convergent and a denominator in (4.21) does not vanish.

We apply formulas (4.21) and (4.22) to calculate the following Dirichlet series $D(\sigma_0, 2, s, p^m) = \sum_{k=1}^{\infty} k^{-s} \sigma_0(k^2 p^m)$ keeping in mind [2] the standard Dirichlet series $D(\sigma_0, 2, s, 1) = \zeta^3(s)/\zeta(2s)$, i.e., $a_1 = 0$. Reduce our problem as follows,

$$D(\sigma_0, 2, s, p^m) = p^{sm/2} \sum_{k=1}^{\infty} \frac{\sigma_0(k^2 p^m)}{(k^2 p^m)^{s/2}} = p^{sm/2} R_{\infty}\left(\frac{\sigma_0(k)}{k^{s/2}}, 2; p^m\right) \frac{\zeta^3(s)}{\zeta(2s)}, \quad (4.23)$$

and calculate the renormalization function in (4.23). According to (4.22) we get

$$p^{sm/2} \mathbf{U}\left(\frac{\sigma_0(k)}{k^{s/2}}, 2; p^m\right) = \frac{p^s}{p^s - 1} \left(m + 1 + \frac{2}{p^s - 1}\right), \quad 1 + \mathbf{V}\left(\frac{\sigma_0(k)}{k^{s/2}}, 2; p\right) = \frac{p^s}{p^s - 1} \left(1 + \frac{2}{p^s - 1}\right)$$

so that following (4.21) and (4.23) we arrive finally at

$$D(\sigma_0, 2, s, p^m) = \frac{(m+1)(p^s - 1) + 2}{p^s + 1} \frac{\zeta^3(s)}{\zeta(2s)}. \quad (4.24)$$

5 RENORMALIZATION OF THE BASIC SUMMATORY FUNCTIONS

In this section we calculate the renormalization function $R_{\infty}(f; p^m)$ for various summatory functions given in Tables 1, 2. For this purpose almost all summatory functions are treated by Theorem 2 and corresponding formulas (2.40) and (2.41) based on calculation of $f(p^r)$. However, in section 5.3 we present another approach, which follows Theorem 1, and calculate the characteristic functions $L_r(f; p^m)$ for the Ramanujan τ function.

5.1 RENORMALIZATION OF SUMMATORY TOTIENT FUNCTIONS

In this section we apply the renormalization approach to summatory totient functions and the Dirichlet series involving the Jordan $J_n(k)$, Euler $\phi(k)$ and Dedekind $\psi(k)$ functions and their combinations. For the two first functions we make use of technical results given in [14]. Discussing the universality classes $\mathbb{B}\{G_1(N); G_2(N)\}$ we will skip hereafter the error term $G_2(N)$.

5.1.1 Euler $\phi(k)$ function

Denote the asymptotics of the summatory functions $F\{k^u \phi^v; N, 1\} = \sum_{k \leq N} k^u \phi^v(k)$ in three different ranges of varying parameters $-\infty < u < \infty$ and $v \in \mathbb{Z}$ which is given in [14],

$$F\{k^u \phi^v; N, 1\} \stackrel{N \rightarrow \infty}{\sim} \begin{cases} A(u, v) N^{u+v+1} & , \quad u + v > -1 , \\ B(u, v) \ln N & , \quad u + v = -1 , \\ C(u, v) & , \quad u + v < -1 , \end{cases} \quad (5.1)$$

such that $A(u, 0) = (u + 1)^{-1}$, $B(-1, 0) = 1$ and $C(u, 0) = \zeta(-u)$.

In the case of arbitrary u and v a lot of expressions for $A(u, v)$, $B(u, v)$ and $C(u, v)$ can be found in [2], [6], [33], [38] and [44]. Here we focus on renormalization functions not specifying the explicit expressions. According to [14], if $v \neq 0$ then $A(u, v)$, $B(u, v)$ and $C(u, v)$ are bounded from above as follows: if $v \in \mathbb{Z}_+$, then

$$0 < A(u, v) \leq \frac{(u + v + 1)^{-1}}{\zeta(v + 1)}, \quad 0 < B(u, v) \leq 1/\zeta(v + 1), \quad 0 < C(u, v) \leq \zeta(-u - v)/\zeta(-u),$$

and if $v \in \mathbb{Z}_-$, then

$$(u + v + 1)^{-1} < A(u, v) \leq 2^{\frac{|v|}{2}} \mathcal{D}_\infty(v, 1)(u + v + 1)^{-1}, \quad 1 < B(u, v) \leq 2^{\frac{|v|}{2}} \mathcal{D}_\infty(v, 1),$$

$$\zeta(-u - v) < C(u, v) \leq 2^{\frac{|v|}{2}} \mathcal{D}_\infty(v, -u - v)\zeta(-u - v), \quad \text{where } \mathcal{D}_\infty(v, s) = \prod_{r=1}^{|v|} \zeta\left(s + \frac{r}{2}\right).$$

Calculate their renormalization functions $R_\infty(k^u \phi^v; p^m)$ according to (2.40) and (2.41),

$$R_\infty(k^u \phi^v; p^m) = p^{m(u+v)+1} \frac{(p-1)^{v-1}}{p^v + (p-1)^{v-1}}, \quad u + v \geq -1, \quad (5.2)$$

$$R_\infty(k^u \phi^v; p^m) = p^{(m-1)(u+v)} \frac{(p-1)^v}{p^{-u} - p^v + (p-1)^v}, \quad u + v < -1.$$

By (5.2) and (5.4) we have an equality $R_\infty(k^u J_1; p^m) = R_\infty(k^u \phi; p^m)$ in all ranges of u .

5.1.2 Jordan $J_v(k)$ and Dedekind $\psi(k)$ functions

Regarding the Jordan function $J_v(k)$, asymptotics of the summatory functions $F\{k^u J_v; N, 1\} = \sum_{k \leq N} k^u J_v(k)$, $v \in \mathbb{Z}_+$, can be given in three different ranges of varying parameters [14],

$$F\{k^u J_v; N, 1\} \stackrel{N \rightarrow \infty}{\sim} \begin{cases} (u + v + 1)^{-1} \frac{N^{u+v+1}}{\zeta(v+1)} & , \quad u + v > -1 , \\ \ln N / \zeta(v + 1) & , \quad u + v = -1 , \\ \zeta(-u - v) / \zeta(-u) & , \quad u + v < -1 , \end{cases} \quad (5.3)$$

and calculate their renormalization function $R_\infty(k^u J_v; p^m)$ in accordance with (2.40) and (2.41)

$$\frac{R_\infty(k^u J_v; p^m)}{p^v - 1} = \frac{p^{m(u+v)+1}}{p^{v+1} - 1}, \quad u + v \geq -1, \quad \frac{R_\infty(k^u J_v; p^m)}{p^v - 1} = \frac{p^{(m-1)(u+v)}}{p^{-u} - 1}, \quad u + v < -1. \quad (5.4)$$

Consider another summatory functions $F\{k^u\psi^v; N, 1\} = \sum_{k \leq N} k^u\psi^v(k)$ in three different ranges of varying parameters $-\infty < u < \infty$ and $v \in \mathbb{Z}$ and note that

$$k^u\psi^v(k) \in \begin{cases} \mathbb{B}\{N^{u+v+1}\} & , \quad u+v > -1, \\ \mathbb{B}\{\ln N\} & , \quad u+v = -1, \\ \mathbb{B}\{N^0\} & , \quad u+v < -1. \end{cases} \quad (5.5)$$

The explicit asymptotics for some summatory functions $F\{k^u\psi^v; N, 1\}$ are given in [40], [44].

Calculate their renormalization functions $R_\infty(k^u\psi^v; p^m)$ according to (2.40) and (2.41),

$$\begin{aligned} R_\infty(k^u\psi^v; p^m) &= p^{m(u+v)+1} \frac{(p+1)^v}{(p+1)^v + (p-1)p^v}, \quad u+v \geq -1, \\ R_\infty(k^u\psi^v; p^m) &= p^{(m-1)(u+v)} \frac{(p+1)^v}{(p+1)^v - p^v + p^{-u}}, \quad u+v < -1. \end{aligned} \quad (5.6)$$

We finish this section with summatory $F\{(\phi/\psi)^v; N, 1\} = \sum_{k \leq N} \phi^v(k)\psi^{-v}(k)$, $v \in \mathbb{Z}$.

The asymptotics $N^{-1}F\{\phi/\psi; N, 1\} \simeq \prod_p(1 - 2/(p(p+1))) \simeq 0.4716$ is known due to [44].

Keeping in mind $(\phi/\psi)^v \in \mathbb{B}\{N\}$ calculate a corresponding renormalization function,

$$R_\infty\left(\left(\frac{\phi}{\psi}\right)^v; p^m\right) = \frac{p(p-1)^{v-1}}{(p-1)^{v-1} + (p+1)^v}, \quad (5.7)$$

which does not depend on m .

5.2 RENORMALIZATION OF SUMMATORY NON-TOTIENT FUNCTIONS

In this section we apply the renormalization approach to summatory functions and Dirichlet series involving divisor $\sigma_a(k)$, prime divisor $\beta(k)$, Piltz $d_n(k)$, abelian group enumeration $\alpha(k)$ functions, Ramanujan sum $C_q(n)$ and some of their combinations.

5.2.1 Divisor function $\sigma_a(k)$ and prime divisor function $\beta(k)$

The divisor function $\sigma_a(k)$ is defined as a sum of the a th powers of the divisors of k . For $k = p^r$ we have $\sigma_a(p^r) = (p^{a(r+1)} - 1) / (p^a - 1)$, $a \neq 0$, and $\sigma_0(p^r) = r + 1$.

- $F\{k^{-s}\sigma_a; N, 1\}$, $a > 0$, $s \geq 1 + a$, $k^{-s}\sigma_a \in \mathbb{B}\{N^0\}$

$$R_\infty\left(\frac{\sigma_a}{k^s}; p^m\right) = \frac{p^{a(m+1)+s} - p^{a(m+1)} - p^s + p^a}{p^{(m+1)s}(p^a - 1)}. \quad (5.8)$$

- $F\{\sigma_a; N, 1\}$, $a > 0$, $\sigma_a \in \mathbb{B}\{N^{a+1}\}$

$$R_\infty(\sigma_a; p^m) = \frac{p^{a(m+1)+1} - p^{am} - p + 1}{p(p^a - 1)}. \quad (5.9)$$

- $F\{\sigma_a; N, 1\}$, $a < 0$, $\sigma_a \in \mathbb{B}\{N\}$

$$R_\infty(\sigma_a; p^m) = \frac{p^{am+1} - p^{am} - p^{1-a} + 1}{p^{1-a}(p^a - 1)}. \quad (5.10)$$

- $F\{\sigma_0^n; N, 1\}$, $\sigma_0^n \in \mathbb{B}\{N(\log N)^{2^n-1}\}$

$$R_\infty(\sigma_0^n; p^m) = \frac{\mathcal{S}(n, p, m+1)}{\mathcal{S}(n, p, 1)}, \quad \mathcal{S}(n, p, t) = \sum_{k=1}^n \binom{n}{k} t^{n-k} Li_{-k}(p^{-1}) + \frac{t^n}{1-p^{-1}}, \quad (5.11)$$

where $Li_{-k}(x)$ is the polylogarithm function defined in (2.14). Substituting $n = 1$ into (5.11) we get $R_\infty(\sigma_0; p^m) = m + 1 - m/p$. According to (5.9) and (5.10), this expression coincides with both limits of $R_\infty(\sigma_a; p^m)$, when $a \rightarrow 0$, for $\sigma_a >$ and $\sigma_a < 0$, respectively.

- $F\{\sigma_0\sigma_a; N, 1\}$, $a > 0$, $\sigma_0\sigma_a \in \mathbb{B}\{N^{a+1} \log N\}$

$$R_\infty(\sigma_0\sigma_a; p^m) = \frac{p^{am+1}(p^{a+1}-1)^2 - p^{a+1}(p-1)^2 + m(p-1)(p^{a+1}-1)(p^{am}(p^{a+1}-1) - p + 1)}{p(p^a-1)(p^{a+2}-1)}$$

such that $R_\infty(\sigma_0\sigma_a; 1) = 1$. Note that $R_\infty(\sigma_0\sigma_a; p^m) \xrightarrow{a \rightarrow 0} R_\infty(\sigma_0^2; p^m)$ according to (5.11).

- $F\{\sigma_a^2; N, 1\}$, $a > 0$, $\sigma_a^2 \in \mathbb{B}\{N^{2a+1}\}$

$$R_\infty(\sigma_a^2; p^m) = \frac{(p-1)(p^{a+1}-1) + p^{am}(p^{1+2a}-1)(p^{am}(p^{1+a}-1) - 2(p-1))}{p(p^a-1)^2(p^{a+1}+1)}, \quad (5.12)$$

that gives $R_\infty(\sigma_a^2; p^m) \xrightarrow{a \rightarrow 0} R_\infty(\sigma_0^2; p^m) = (p + (m(p-1) + p)^2)/(p(p+1))$ according to (5.11).

- $F\{1/\sigma_0; N, 1\}$, $1/\sigma_0 \in \mathbb{B}\{N/\sqrt{\log N}\}$

$$R_\infty\left(\frac{1}{\sigma_0}; p^m\right) = -\frac{{}_2F_1(m+1, 1; m+2; p^{-1})}{(m+1)p \ln(1-p^{-1})}, \quad (5.13)$$

where ${}_2F_1(a, b; c; z)$ denotes a generalized hypergeometric function [12]. By (5.13) we get for $m = 1$

$$R_\infty\left(\frac{1}{\sigma_0}; p\right) = p + \frac{1}{\ln(1-p^{-1})}, \quad R_\infty\left(\frac{1}{\sigma_0}; p\right) \xrightarrow{p \rightarrow \infty} \frac{1}{2}, \quad R_\infty\left(\frac{1}{\sigma_0}; p\right) \xrightarrow{p \rightarrow 1} 1. \quad (5.14)$$

Expanding an expression (5.13) as an infinite series $\sum_{r=0}^{\infty} L_r(1/\sigma_0; p^m) p^{-r}$ in accordance with Theorem 1, one can calculate $L_r(1/\sigma_0; p^m)$ and verify that for $0 \leq r \leq 3$ they coincide with those given in (1.17).

- $F\{\sigma_1/\sigma_0; N, 1\}$, $\sigma_1/\sigma_0 \in \mathbb{B}\{N^2/\sqrt{\log N}\}$

$$R_\infty\left(\frac{\sigma_1}{\sigma_0}; p^m\right) = \frac{p^{m+1} {}_2F_1(m+1, 1; m+2; p^{-1}) - {}_2F_1(m+1, 1; m+2; p^{-2})}{(m+1)p^2 \log(1+p^{-1})}, \quad (5.15)$$

that gives for $m = 1$

$$R_\infty\left(\frac{\sigma_1}{\sigma_0}; p\right) = p^2 - \frac{p-1}{\log(1+p^{-1})}, \quad \frac{1}{p} R_\infty\left(\frac{\sigma_1}{\sigma_0}; p\right) \xrightarrow{p \rightarrow \infty} \frac{1}{2}. \quad (5.16)$$

- $F\{\sigma_1/\phi; N, 1\}, F\{\sigma_1/\psi; N, 1\}, \quad \sigma_1/\phi, \sigma_1/\psi \in \mathbb{B}\{N\}, m \geq 1.$

$$R_\infty\left(\frac{\sigma_1}{\phi}; p^m\right) = \frac{p^3(1+p-p^{-m})}{p^4-p^3+p^2+p-1}, \quad R_\infty\left(\frac{\sigma_1}{\psi}; p^m\right) = \frac{p^3(1+p-p^{-m})}{p^4+p^3-p^2-p+1}. \quad (5.17)$$

The prime divisor function $\beta(k)$ is defined by formula $\beta(p^{a_1} \cdots p^{a_n}) = a_1 \cdots a_n$.

- $F\{\beta; N, 1\}, \beta \in \mathbb{B}\{N\}$

$$R_\infty(\beta; p^m) = p \frac{m(p-1)+1}{p^2-p+1}. \quad (5.18)$$

Let us note a curious consequence of (5.18) when $m = p$: $R_\infty(\beta; p^p) = p$.

- $F\{k^{-s}\beta; N, 1\}, \beta \in \mathbb{B}\{N^0\}$

$$R_\infty\left(\frac{\beta}{k^s}; p^m\right) = p^{(1-m)s} \frac{m(p^s-1)+1}{p^{2s}-p^s+1}. \quad (5.19)$$

A generalized summatory function $F\{\beta^n; N, 1\}$ was considered in [27] such that $\mathcal{F}(\beta^n) = \prod_{p \geq 2} \left(1 + \sum_{j=2}^{\infty} [j^n - (j-1)^k] p^{-j}\right)$, e.g., $\mathcal{F}(\beta) = \prod_{p \geq 2} [1 - 1/(p(p-1))] = \zeta(2)\zeta(3)/\zeta(6)$.

- $F\{\beta^n; N, 1\}, \beta^n \in \mathbb{B}\{N\}$

$$R_\infty(\beta^n; p^m) = \frac{\mathcal{S}(n, p, m)}{1 + Li_{-n}(p^{-1})}, \quad (5.20)$$

where $\mathcal{S}(n, p, m)$ is defined in (5.11).

5.2.2 Piltz function $d_n(k)$ and the sum of two squares function $r_2(k)$

The Piltz function $d_n(k)$ is defined as a number of ways to write the positive integer k as a product of n (positive integer) factors. For $k = p^r$ we have $d_n(p^r) = \binom{n+r-1}{r}$. By definition, it holds $d_1(k) = 1$ and $d_2(k) = \sigma_0(k)$.

- $F\{k^{-s}d_n; N, 1\}, d_n \in \mathbb{B}\{N^0\}$

$$R_\infty\left(\frac{d_n}{k^s}; p^m\right) = p^{-sm} \binom{n+m-1}{m} {}_2F_1(m, 1-n; m+1; p^{-s}). \quad (5.21)$$

- $F\{d_n; N, 1\}, d_n \in \mathbb{B}\{N(\log N)^{n-1}\}$

$$R_\infty(d_n; p^m) = \binom{n+m-1}{m} {}_2F_1(m, 1-n; m+1; p^{-1}). \quad (5.22)$$

Note that $R_\infty(d_2; p^m)$ is coincided with $R_\infty(\sigma_0; p^m)$ given in section 5.2.1.

- $F\{d_n^2; N, 1\}, d_n^2 \in \mathbb{B}\{N(\log N)^{n^2-1}\}$

$$R_\infty(d_n^2; p^m) = \binom{n+m-1}{m}^2 \frac{{}_3F_2(1, m+n, m+n; m+1, m+1; p^{-1})}{{}_2F_1(n, n; 1; p^{-1})}, \quad (5.23)$$

and $R_\infty(d_2^2; p^m) = R_\infty(\sigma_0^2; p^m)$ in accordance with (5.11).

- $F\{1/d_n; N, 1\}, 1/d_n \in \mathbb{B}\{N(\log N)^{1/n-1}\}$

$$R_\infty\left(\frac{1}{d_n}; p^m\right) = np \binom{n+m-1}{m}^{-1} \frac{{}_2F_1(1, m+1; m+n; p^{-1})}{n p + {}_2F_1(2, 1; n+1; p^{-1})}, \quad (5.24)$$

such that $R_\infty(d_2^{-1}; p^m) = R_\infty(\sigma_0^{-1}; p^m)$ in accordance with (5.13).

The number of representations of k by two squares, allowing zeros and distinguishing signs and order, is denoted by $r_2(k)$. If $k = p^r$ then,

$$r_2(p^r) = \begin{cases} 4(r+1) & , \quad p = 1 \pmod{4}, \\ 4 & , \quad p = 2 \end{cases} \quad \text{and} \quad r_2(p^r) = \begin{cases} 4 & , \quad p = 3 \pmod{4}, 2 \mid r \\ 0 & , \quad p = 3 \pmod{4}, 2 \nmid r \end{cases}$$

- $F\{r_2; N, 1\}, r_2 \in \mathbb{B}\{N\}$

$$R_\infty(r_2; p^m) = \begin{cases} m+1-m/p & , \quad p = 1 \pmod{4}, \\ 1 & , \quad p = 2, \end{cases} \quad R_\infty(r_2; p^m) = \begin{cases} 1 & , \quad p = 3 \pmod{4}, 2 \mid m \\ 1/p & , \quad p = 3 \pmod{4}, 2 \nmid m \end{cases}$$

- $F\{r_2/k; N, 1\}, r_2/k \in \mathbb{B}\{\log N\}$

$$R_\infty\left(\frac{r_2}{k}; p^m\right) = p^{-m} R_\infty(r_2; p^m) \quad (5.25)$$

5.2.3 Ramanujan sum $C_q(n)$ and Abelian group enumeration function $\alpha(k)$

Ramanujan's sum $C_q(n)$, $q, n \geq 1$, is a multiplicative arithmetic function which is defined by formula $C_q(n) = \sum_{a=1}^q \exp(2\pi i n a/q)$, $(a, q) = 1$ such that $C_{q_1 q_2}(n) = C_{q_1}(n) C_{q_2}(n)$ if $(q_1, q_2) = 1$ and $C_1(n) = 1$. If $q = p^r$ then

$$\begin{aligned} C_{p^r}(n) &= 0, \text{ if } p^{r-1} \nmid n; \quad C_{p^r}(n) = -p^{r-1}, \text{ if } p^{r-1} \mid n, p^r \nmid n; \quad C_{p^r}(n) = \phi(r), \text{ if } p^r \mid n, \\ C_p(n) &= -1, \text{ if } p \nmid n; \quad C_{p^r}(n) = 0, \text{ if } p \nmid n \text{ and } r \geq 2. \end{aligned} \quad (5.26)$$

Consider the Dirichlet summatory function for $C_q(n)$: $F\{C_q(n)q^{-s}; \infty, 1, s\} = \sum_{q=1}^{\infty} C_q(n)q^{-s}$, $s > 1$, where n is kept constant. It is convergent to $n^{-s+1}\sigma_{s-1}(n)\zeta^{-1}(s)$. We find a rescaled summatory function $F\{C_q(n)q^{-s}; \infty, p^m, s\} = \sum_{q=1}^{\infty} C_{p^m q}(n)q^{-s}$.

Let a number n is such that $p^a \mid n$ but $p^{a+1} \nmid n$, $a \in \mathbb{Z}_+$, and $a \geq m \geq 1$, then

$$\mathbf{U}\left(\frac{C_q(n)}{q^s}; p^m\right) = p^{-ms} \left(\sum_{r=m}^a \frac{\phi(p^r)}{p^{sr}} - \frac{p^a}{p^{s(a+1)}} \right), \quad \mathbf{V}\left(\frac{C_q(n)}{q^s}; p^m\right) = \sum_{r=1}^a \frac{\phi(p^r)}{p^{sr}} - \frac{p^a}{p^{s(a+1)}} \quad (5.27)$$

that gives

$$F\left\{\frac{C_q(n)}{q^s}; \infty, p^m, s\right\} = \frac{p^s - 1 + p^{(s-1)(a+2-m)} - 1}{(p^s - 1)(p^{(s-1)(a+1)})} F\left\{\frac{C_q(n)}{q^s}; \infty, 1, s\right\}. \quad (5.28)$$

If $a \geq m = 0$ then by (5.27) we have $\mathbf{U}(C_q(n)q^{-s}; 1) = 1 + \mathbf{V}(C_q(n)q^{-s}; 1)$, and therefore $F\{C_q(n)q^{-s}; \infty, p^m, s\} \xrightarrow{m \rightarrow 0} F\{C_q(n)q^{-s}; \infty, 1, s\}$.

If $m \geq 2$ and a number n is not divided by p , i.e., $a = 0$, then $F\{C_q(n)q^{-s}; \infty, p^m, s\} = 0$.

Finally, if $m = 1$ and $a = 0$, then $F\{C_q(n)q^{-s}; \infty, p, s\} = F\{C_q(n)q^{-s}; \infty, 1, s\} / (p^{-s} - 1)$.

Abelian group enumeration function $\alpha(k)$ accounts for the number of (isomorphism classes of) commutative groups of order k . By definition it satisfies $\alpha(p^r) = \mathcal{P}(r)$, where $\mathcal{P}(r)$ denotes an unrestricted partition function [1] and $\mathcal{P}(0) = 1$.

- $F\{\alpha; N, 1\}$, $\alpha \in \mathbb{B}\{N\}$

$$R_\infty(\alpha; p^m) = p^m \left(1 - Q(p^{-1}) \sum_{k=0}^{m-1} \mathcal{P}(k) p^{-k} \right), \quad Q(x) = \prod_{j=1}^{\infty} (1 - x^j) \quad |x| \leq 1. \quad (5.29)$$

- $F\{1/\alpha; N, 1\}$, $1/\alpha \in \mathbb{B}\{N\}$

$$R_\infty\left(\frac{1}{\alpha}; p^m\right) = p^m \frac{\mathcal{T}(p, m)}{\mathcal{T}(p, 0)}, \quad \mathcal{T}(p, t) = \sum_{k=t}^{\infty} \frac{p^{-k}}{\mathcal{P}(k)}. \quad (5.30)$$

5.3 RENORMALIZATION OF SUMMATORIES ASSOCIATED WITH RAMANUJAN'S τ FUNCTION

The Ramanujan τ function is a multiplicative arithmetic function which is mostly known due to its generating function, $\sum_{k=1}^{\infty} \tau(k)x^k = x \prod_{k=1}^{\infty} (1 - x^k)^{24}$, $|x| < 1$. For our purpose to calculate the renormalization function for any summatory function $F\{f(\tau, k); N, 1\}$ with $f(\tau, k)$ involving the τ function, it is important to know its recursive relation for $k = p^r$,

$$\tau(p^r) = \sum_{j=0}^{\lfloor r/2 \rfloor} (-1)^j \binom{r-j}{r-2j} p^{11j} \tau^{r-2j}(p). \quad (5.31)$$

Then, making use of formulas (2.41) and (2.42) we can arrive at $R_\infty(f(\tau, k); p^m)$ due to the finite computational procedure. However, the representation (5.31) is too difficult to make worth, so we choose another way to find $R_\infty(f(\tau, k); p^m)$, namely, by Theorem 1 and by calculating the characteristic functions $L_r(\tau; p^m)$. Start with identity for τ function [31]

$$\tau(p^{r+1}) = \tau(p)\tau(p^r) - p^{11}\tau(p^{r-1}). \quad (5.32)$$

The following Proposition is based on recursion (1.14) and the last identity.

Proposition 2

$$L_0(\tau; p^m) = \tau(p^m), \quad L_1(\tau; p^m) = -p^{11}\tau(p^{m-1}), \quad L_r(\tau; p^m) = 0, \quad r \geq 2. \quad (5.33)$$

Proof Calculating the four first expressions of $L_r(\tau; p^m)$ one can verify that (5.33) holds, i.e., $L_2(\tau; p^m) = L_3(\tau; p^m) = 0$. Prove by induction that $L_r(\tau; p^m) = 0$, $r \geq 2$.

Indeed, let $L_q(\tau; p^m) = 0$ for $2 \leq q \leq r$, then keeping in mind (1.14) and (5.32) write this equality in another representation,

$$L_q(\tau; p^m) = \tau(p^{m+q}) - \tau(p^m)\tau(p^q) + p^{11}\tau(p^{m-1})\tau(p^{q-1}) = 0, \quad 2 \leq q \leq r. \quad (5.34)$$

Making use of (5.32) and (5.34) write the next term $L_{r+1}(\tau; p^m)$,

$$L_{r+1}(\tau; p^m) = \tau(p^{m+r+1}) - \tau(p^m)\tau(p^{r+1}) + p^{11}\tau(p^{m-1})\tau(p^r),$$

and calculate a difference,

$$\begin{aligned} L_{r+1}(\tau; p^m) - \tau(p)L_r(\tau; p^m) &= [\tau(p^{m+r+1}) - \tau(p)\tau(p^{m+r})] - \tau(p^m)[\tau(p^{r+1}) - \tau(p)\tau(p^r)] \\ &\quad + p^{11}\tau(p^{m-1})[\tau(p^r) - \tau(p)\tau(p^{r-1})]. \end{aligned} \quad (5.35)$$

By identity (5.32) the r.h.s. in equality (5.35) can be reduced as follows,

$$L_{r+1}(\tau; p^m) - \tau(p)L_r(\tau; p^m) = -p^{11}[\tau(p^{m+r-1}) - \tau(p^m)\tau(p^{r-1}) - p^{11}\tau(p^{m-1})\tau(p^{r-2})].$$

By comparison with (5.34) one can recognize the function $L_{r-1}(\tau; p^m)$ in the brackets of the last expression. Then, combining this fact with (5.35) and assumption (5.34) we get

$$L_{r+1}(\tau; p^m) = \tau(p)L_r(\tau; p^m) - p^{11}L_{r-1}(\tau; p^m) = 0. \quad (5.36)$$

Thus, proof is finished. $\square$

An important spinoff arising from Proposition 2 is the identity (5.34) which does generalize an identity (5.32).

Corollary 2 *The Ramanujan τ function satisfies an identity,*

$$\tau(p^{m+n}) = \tau(p^m)\tau(p^n) - p^{11}\tau(p^{m-1})\tau(p^{n-1}), \quad m, n \geq 1. \quad (5.37)$$

The last statement implies two inequalities which could easily be verified. The 1st inequality looks quite trivial, $\tau(p^{2n}) < \tau^2(p^n)$. Regarding the 2nd inequality, let p_* and n_* be chosen in such a way that $\tau(p_*^{2n_*}) < 0$, e.g., $\tau(2^2), \tau(3^2), \tau(5^2), \tau(7^2) < 0$, $\tau(5^4), \tau(11^4) < 0$ etc. Then the following inequality holds,

$$|\tau(p_*^{2n_*})| < p_*^{11}\tau^2(p_*^{n_*-1}). \quad (5.38)$$

In the case $n_* = 2$ let us combine (5.38) with Deligne's inequality $|\tau(p)| < 2p^{11/2}$ for the Ramanujan τ functions [7] and get,

$$|\tau(p_*)^4| < 4p_*^{22}. \quad (5.39)$$

Note that the upper bound in (5.39) is stronger than the bound which came by combining (5.32) and Deligne's inequality for arbitrary prime p . Indeed,

$$\tau(p^4) = \tau^4(p) - 3p^{11}\tau^2(p) + p^{22} < 17p^{22} - 3p^{11}\tau^2(p). \quad (5.40)$$

However, $4p^{22} < 17p^{22} - 3p^{11}\tau^2(p)$ that follows by Deligne's inequality, $17p^{22} - 3p^{11}\tau^2(p) > 5p^{22}$.

Straightforward numerical calculations show that the inequality (5.39) holds also for the first 474 primes irrespectively whether the requirement $\tau(p^4) < 0$ is holding,

$$p_{475} = 3371, \quad \frac{\tau(3371^4)}{4 \cdot 3371^{22}} \simeq 1.0119.$$

5.3.1 Renormalization of the Ramanujan τ Dirichlet series

Recalling the Ramanujan conjecture on τ function, $\tau(N) = \mathcal{O}(N^{11/2+\varepsilon})$ proved by Deligne, consider the τ Dirichlet series $F\{\tau \cdot k^{-s}; \infty, p^m\} = \sum_{k=1}^{\infty} \tau(k) k^{-s}$, $\tau \cdot k^{-s} \in \mathbb{B}\{N^0\}$, which converges absolutely if $s > 13/2$. By Theorem 1 and relationship (4.14) between characteristic functions for $\tau(k)$ and $\tau(k) k^{-s}$ the renormalization function $R_{\infty}(\tau \cdot k^{-s}; p^m)$ reads

$$R_{\infty}(\tau \cdot k^{-s}, p^m) = \sum_{r=0}^{\infty} L_r(\tau \cdot k^{-s}; p^m) = \sum_{r=0}^{\infty} L_r(\tau; p^m) p^{-s(r+m)}. \quad (5.41)$$

Applying Proposition 2 to (5.41) we obtain,

$$R_{\infty}(\tau \cdot k^{-s}, p^m) = p^{-sm} \left(\tau(p^m) - \frac{\tau(p^{m-1})}{p^{s-11}} \right).$$

Making use of relation between renormalization function $R_{\infty}(f \cdot k^{-s}; p^m)$ and a ratio $\mathfrak{D}(f; p^m, s)$ between scaled and unscaled Dirichlet series, given in section 4.2, we get finally,

$$\sum_{k=1}^{\infty} \frac{\tau(kp^m)}{k^s} = \left(\tau(p^m) - \frac{\tau(p^{m-1})}{p^{s-11}} \right) \sum_{k=1}^{\infty} \frac{\tau(k)}{k^s}. \quad (5.42)$$

Formula (5.42) gives rise to several special cases, e.g.,

$$\sum_{k=1}^{\infty} \frac{\tau(kp)}{k^s} = (\tau(p) - p^{11-s}) \sum_{k=1}^{\infty} \frac{\tau(k)}{k^s}, \quad \sum_{n=0}^m \sum_{k=1}^{\infty} \frac{\tau(kp^n)}{k^{11}} = \tau(p^m) \sum_{k=1}^{\infty} \frac{\tau(k)}{k^{11}}.$$

5.3.2 Renormalization functions $R_\infty(\tau^2, p^m)$ and $R_\infty(\tau^2 \cdot k^{-25/2}, p^m)$

In this section we renormalize the summatory functions $F\{\tau^2; N, p^m\}$ and $F\{\tau^2 \cdot k^{-25/2}; N, p^m\}$ given in Tables 1 and 2. For this purpose we start by calculating the characteristic functions $L_r(\tau^2; p^m)$. In contrast to $L_r(\tau; p^m)$ described in Proposition 2 the present case is not so simple but still allows to find the general expressions.

Proposition 3

$$\begin{aligned} L_0(\tau^2; p^m) &= \tau^2(p^m), \quad L_1(\tau^2; p^m) = p^{22}\tau^2(p^{m-1}) - 2p^{11}\tau(p)\tau(p^{m-1})\tau(p^m), \\ L_r(\tau^2; p^m) &= 2(-p^{11})^r \tau(p)\tau(p^{m-1})\tau(p^m), \quad r \geq 2. \end{aligned} \quad (5.43)$$

Proof Prove Proposition by induction. First, calculating the five first expressions of $L_r(\tau^2; p^m)$ one can verify that (5.43) holds. Let Proposition holds for $2 \leq q \leq r$, then prove that

$$L_{r+1}(\tau^2; p^m) = 2(-p^{11})^{r+1} \tau(p)\tau(p^{m-1})\tau(p^m).$$

Keeping in mind (1.14) calculate $L_{r+1}(\tau^2; p^m)$,

$$\begin{aligned} L_{r+1}(\tau^2; p^m) &= \tau^2(p^{m+r+1}) - \tau^2(p^{r+1})\tau^2(p^m) - \tau^2(p^r)p^{11}\tau(p^{m-1})[p^{11}\tau(p^{m-1}) - 2\tau(p)\tau(p^m)] \\ &\quad - 2\tau(p)\tau(p^{m-1})\tau(p^m)p^{22}\left[\tau^2(p^{r-1}) - \tau^2(p^{r-2})p^{11} + \dots + (-1)^{r+1}\tau^2(p)p^{11(r-2)}\right]. \end{aligned}$$

By (5.32) the four first terms in the above equality are reduced up to a single term,

$$\begin{aligned} \tau^2(p^{m+r+1}) - \tau^2(p^{r+1})\tau^2(p^m) - \tau^2(p^r)p^{11}\tau(p^{m-1})[p^{11}\tau(p^{m-1}) - 2\tau(p)\tau(p^m)] &= \\ 2\tau(p^m)\tau(p^{m-1})\tau(p^r)p^{11}[\tau(p^r)\tau(p) - \tau(p^{r+1})] &= 2\tau(p^m)\tau(p^{m-1})\tau(p^r)\tau(p^{r-1})p^{22}, \end{aligned}$$

which simplifies further calculations,

$$\begin{aligned} L_{r+1}(\tau^2; p^m) &= 2\tau(p^m)\tau(p^{m-1})p^{22}A_{r-1}, \quad \text{where} \\ A_{r-1} &= \tau(p^{r-1})[\tau(p^r) - \tau(p)\tau(p^{r-1})] + \tau(p)\tau^2(p^{r-2})p^{11} - \dots \pm \tau^3(p)p^{11(r-2)}. \end{aligned} \quad (5.44)$$

Performing calculations in curl brackets by (5.32), we obtain

$$\begin{aligned} L_{r+1}(\tau^2; p^m) &= -2\tau(p^m)\tau(p^{m-1})p^{33}A_{r-2}, \quad \text{where} \\ A_{r-2} &= \tau(p^{r-2})[\tau(p^{r-1}) - \tau(p)\tau(p^{r-2})] + \tau(p)\tau^2(p^{r-3})p^{11} - \dots \pm \tau^3(p)p^{11(r-3)}. \end{aligned} \quad (5.45)$$

By comparison (5.44) and (5.45) and continuing to contract the terms in curl brackets, we get

$$L_{r+1}(\tau^2; p^m) = (-1)^r 2\tau(p^m)\tau(p^{m-1})p^{11r}A_1, \quad A_1 = \tau(p)[\tau(p^2) - \tau^2(p)] = -\tau(p)p^{11}.$$

Thus, Proposition is proven. $\square$

Calculate the renormalization function $R_\infty(\tau^2, p^m)$. In accordance with Table 1 we obtain $\tau^2 \in \mathbb{B}\{N^{12}\}$, i.e., $a_1 = 12$. Then by Theorem 1 and Proposition 3 we get

$$R_\infty(\tau^2, p^m) = \sum_{r=0}^{\infty} \frac{L_r(f; p^m)}{p^{12r}} = \tau^2(p^m) + p^{10} \tau^2(p^{m-1}) - \frac{2\tau(p)}{p+1} \tau(p^{m-1}) \tau(p^m), \quad (5.46)$$

such that $R_\infty(\tau^2, 1) = 1$. In the case $m = 1$, we find $R_\infty(\tau^2, p) = p^{10} + \tau^2(p) \cdot (p-1)/(p+1)$.

Finally, calculate the renormalization function $R_\infty(\tau^2 \cdot k^{-25/2}, p^m)$ such that in accordance with Table 2 we have $\tau^2 \cdot k^{-25/2} \in \mathbb{B}\{N^0\}$. Making use of relationship (4.14) between characteristic functions $L_r(f \cdot k^{-s}; p^m)$ and $L_r(f; p^m)$ build the renormalization function as it was done in formula (5.41) for the Ramanujan τ Dirichlet series,

$$R_\infty\left(\frac{\tau^2}{k^{25/2}}, p^m\right) = \sum_{r=0}^{\infty} L_r\left(\frac{\tau^2}{k^{25/2}}; p^m\right) = p^{-25m/2} \sum_{r=0}^{\infty} \frac{L_r(\tau^2; p^m)}{p^{25r/2}}.$$

By Proposition 3 and the above formula we get

$$p^{25m/2} R_\infty\left(\frac{\tau^2}{k^{25/2}}, p^m\right) = \tau^2(p^m) + p^{19/2} \tau^2(p^{m-1}) - \frac{2\tau(p)}{p^{3/2}+1} \tau(p^{m-1}) \tau(p^m). \quad (5.47)$$

6 NUMERICAL VERIFICATION

In this section we verify the renormalization approach for summation of multiplicative arithmetic functions with scaled summation variable by numerical calculations. Consider a relative deviation $\rho(f; N, p^m)$ between rescaled $F\{f; N, p^m\}$ and nonscaled $F\{f; N, 1\}$ summatory functions,

$$\rho(f; N, p^m) = R_\infty^{-1}(f; p^m) \cdot \frac{F\{f; N, p^m\}}{F\{f; N, 1\}} - 1, \quad (6.1)$$


where the renormalization function $R_\infty(f; p^m)$ is calculated according to Theorem 1. In Figures 1, 2 and 3 we present plots of $\rho(f; N, p^m)$ for six different arithmetic functions. These Figures show that formulas for renormalization functions work with high precision.

ACKNOWLEDGEMENT


The useful discussions with A. Juhasz, A. Mann and Z. Rudnick are highly appreciated.

REFERENCES

- [1] G. E. Andrews, *The Theory of Partitions*, Encyclopedia of mathematics and its applications, **2**, Addison-Wesley, Reading, MA, 1976




(a)




(b)

Figure 1: Plots of $\rho(\phi; N, 11)$ (a) and $\rho(\sigma_1; N, 11)$ (b) in the range $2 \cdot 10^4 \leq N \leq 10^6$.




(a)




(b)

Figure 2: Plots of $\rho(\sigma_{-1}; N, 11)$ (a) and $\rho(|\mu|; N, 11)$ (b) in the range $2 \cdot 10^4 \leq N \leq 10^6$.



(a)



(b)

Figure 3: Plots of $\rho(\tau \cdot k^{-7}; N, 3)$ in the range $5 \cdot 10^3 \leq N \leq 10^5$ (a) and $\rho(\tau^2; N, 3)$ (b) in the range $5 \cdot 10^3 \leq N \leq 3 \cdot 10^5$.

[2] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, N.Y, 1995, 4th ed.

- [3] P. T. Bateman, P. Erdős, C. Pomerance and E. G. Strauss, *The Arithmetic Mean of the Divisors of an Integer*, Analytic Number Theory, Proceed. 1980 Phil. Confer., ed. M. I. Knopp, Lect. Notes in Math., **899**, Springer, 1981, pp. 197-220
- [4] U. Balakrishnana and Y. -F. S. Pétermann, *Asymptotic Estimates for a Class of Summatory Functions*, J. Number Theory, **70**, 1-36 (1998)
- [5] K. A. Broughan, *Restricted divisor sums*, Acta Arith., **101**, 105-114 (2002)
- [6] S. D. Chlowla, *An Order Result Involving Euler's ϕ - Function*, J. Indian Math. Soc., **18**, 138-141 (1927)
- [7] P. Deligne, *La Conjecture de Weil. I*. Inst. Hautes Études Sci. Publ. Math., **43**, 273-307 (1974)
- [8] R. Diaz and S. Robins, *The Ehrhart Polynomial of a Lattice Polytope*, Ann. Math., **145**, # 3, 503-518 (1997)
- [9] L. E. Dickson, *History of the Theory of Numbers*, **1**, Chelsea Pub.Co, N.Y, 1952
- [10] T. Dokchitser, *Computing special values of motivic L-functions*, Experimental. Math., **13**, 137-149 (2004)
- [11] P. D. T. A. Elliott, *Multiplicative Functions and Ramanujan's τ -Function*, J. Austral. Math. Soc., **A 30**, 461-468 (1980/81)
- [12] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions*, Vol. 1. New York: Krieger, 1981.
- [13] P. Erdős and G. Szekeres, *Über die Anzahl der Abelschen Gruppen gegebener Ordnung und über ein Verwandtes Zahlentheoretisches Problem*, Acta Sci. Math. (Szeged) **7**, 95-102 (1935)
- [14] L. G. Fel, *On Summatory Totient Functions*, preprint, (2008), <http://arXiv.org/abs/math.NT/0802.0619>
- [15] C. Ferreira, J. L. López, *Asymptotic Expansions of the Hurwitz-Lerch zeta Function*, J. Math. Anal. Appl. **298**, 210-242 (2004)
- [16] S. R. Finch, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 106, 104-110
- [17] D. Goldfeld and P. Sarnak, *Sums of Kloosterman Sums*, Invent. math., **71**, 243-250 (1983)

- [18] L. Greenberg, M. Newman, *Some Results on Solvable Groups*, Arch. Math., **21**, 349 (1970)
- [19] E. Grosswald, *The Average Order of Arithmetic Function*, Duke Math. J. **23** 41-44 (1956)
- [20] G. H. Hardy, *On Dirichlet's Divisor Problem*, Proc. Lond. Math. Soc., 2, **15**, # 1251 (1915)
- [21] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*,
Oxford : Clarendon Press, 1979, 5th ed.
- [22] G. Higman, *Enumerating of p -groups*, Proc. Lond. Math. Soc., **10**, 24-30 (1960)
- [23] E. M. Horadam, *Ramanujan's Sum and Nagell's Totient Function for Arithmetic Semigroups*, Math. Scand., **22**, 269-281 (1968)
- [24] A. Ivić, *On the Asymptotic Formulae for Some Functions Connected with Powers of the Zeta-Function*, Mat. Vestnik (Belgrade), 1 (14), **29**, 79-90 (1977).
- [25] V. Kalinka, *A Variant of the Divisor Problem with a Large Number of Components*, Lith. Math. J., **14**, 442-448 (1974)
- [26] D. G. Kendall and R. A. Rankin, *On the Number of Abelian Groups of Given Order*, Quart. J. Math. Oxford Ser. **18**, 197-208 (1947)
- [27] J. Knopfmacher, *A Prime-Divisor Function*, Proc. Amer. Math. Soc., **40**, 373-377 (1973)
- [28] J.-M. de Koninck and A. Ivić, *Topics in Arithmetical Functions : Asymptotic Formulae for Sums of Reciprocals of Arithmetical Functions and Related Results*,
Amsterdam : North-Holland Pub. Co., 1980, Chapt. 1
- [29] E. Krätzel, *Lattice Points*, Berlin - Dordrecht: Kluwer, 1988, pp. 142-143
- [30] E. Landau, *Über die Zahlentheoretische Function $\phi(n)$ und ihre Beziehung zum Goldbachschen Satz*, Nachr. Königlichen Ges. Wiss. Göttingen, Math.-Phys. Klasse, 177-186, 1900.
- [31] D. H. Lehmer, *Ramanujan's Function $\tau(n)$* , Duke Math. J., **10**, 483-492 (1943)
- [32] A. Lubotzky and D. Segal, *Subgroup growth*, Basel : Birkhauser, 2003, p. 22
- [33] W. G. Nowak, *On an Error Term Involving the Totient Function*, Indian J. Pure Appl. Math., **20**, 537-542 (1989)
- [34] S. Ramanujan, *Some Formulae in the Analytic Theory of Numbers*, Messenger of Math., **45**, 81-84 (1915)

- [35] R. A. Rankin, *Sums of Powers of Cusp Form Coefficients*,
Math. Annalen, **263**, 227-236 (1983)
- [36] M. P. F. du Sautoy, *Counting p -groups and Nilpotent Groups*,
Publications Mathématiques de l'IHÉS, 92 (2000), p. 63-112
- [37] J. Sandór, *On the Arithmetical Functions $d_n(k)$ and $d_n^*(k)$* ,
Portugaliae Math., **53**, 107-115 (1996)
- [38] J. Sandór, D. S. Mitrinović and B. Crstici, *Handbook of Number Theory I*,
Springer, Dordrecht, The Netherlands, 2006, 2nd ed
- [39] A. Selberg, *Note on the paper by L. G. Sathe*,
J. Indian Math. Soc. (Old Ser.), **28**, 83-87 (1954)
- [40] V. Sita Ramaiah and D. Suryanarayana, *Sums of Reciprocals of Some
Multiplicative Functions*, Math. J. Okayama Univ., **21**, 155-164 (1979)
- [41] R. Sitaramachandra Rao, *On an Error Term of Landau. II*,
Rocky Mountain J. Math. **15**, 579-588 (1985)
- [42] R. A. Smith, *An Error Term of Ramanujan*, J. Number Theory, **2**, 91-96 (1970)
- [43] M. Spivak, *Calculus*, Berkeley, Calif.: Publish or Perish, c1980.
- [44] D. Suryanarayana, *On Some Asymptotic Formulae of S. Wigert*,
Indian J. Math., **24**, 81-98 (1982)
- [45] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*,
Cambridge : Cambridge University Press, c1995
- [46] A. Walfisz, *Weylsche Exponentialsummen in der Neueren Zahlentheorie*,
VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.
- [47] D. R. Ward, *Some Series Involving Euler Function*, J. London Math. Soc., **2**, 210-214 (1927)
- [48] B. M. Wilson, *Proofs of Some Formulae Enunciated by Ramanujan*,
Proc. London Math. Soc., **21**, pp. 235-255 (1923)